

4-7-2015

# Is there specificity in a defensive mutualism against soil versus lab nematodes, *Dictyostelium discoideum* farmers and their bacteria?

Boahemaa Adu-Oppong,


David C. Queller

*Washington University in St Louis*, [queller@WUSTL.EDU](mailto:queller@WUSTL.EDU)

Joan E. Strassmann

*Washington University in St Louis*, [strassmann@WUSTL.EDU](mailto:strassmann@WUSTL.EDU)

Follow this and additional works at: [http://openscholarship.wustl.edu/bio\\_facpubs](http://openscholarship.wustl.edu/bio_facpubs)

 Part of the [Behavior and Ethology Commons](#), [Biology Commons](#), and the [Environmental Microbiology and Microbial Ecology Commons](#)

---

## Recommended Citation

Adu-Oppong,, Boahemaa; Queller, David C.; and Strassmann, Joan E., "Is there specificity in a defensive mutualism against soil versus lab nematodes, *Dictyostelium discoideum* farmers and their bacteria?" (2015). *Biology Faculty Publications & Presentations*. Paper 62. [http://openscholarship.wustl.edu/bio\\_facpubs/62](http://openscholarship.wustl.edu/bio_facpubs/62)

This Article is brought to you for free and open access by the Biology at Washington University Open Scholarship. It has been accepted for inclusion in Biology Faculty Publications & Presentations by an authorized administrator of Washington University Open Scholarship. For more information, please contact [digital@wumail.wustl.edu](mailto:digital@wumail.wustl.edu).

## Is there specificity in a defensive mutualism against soil versus laboratory nematodes, *Dictyostelium discoideum* farmers and their bacteria?

Boahemaa Adu-Oppong, David C. Queller and Joan E. Strassmann

Department of Biology, Washington University in St. Louis, St. Louis, Missouri, USA

---

### ABSTRACT

**Background:** The social amoeba *Dictyostelium discoideum* is a soil-dwelling microbe, which lives most of its life cycle in the vegetative stage as a predator of bacteria and as prey for nematodes. When bacteria are sparse, amoebae aggregate into a multicellular fruiting body. Some clones of *D. discoideum* have agriculture (Brock *et al.*, 2011). They carry bacteria through the social stage, eat them prudently, and use some bacteria as defence against non-farming *D. discoideum* competitors. *Caenorhabditis elegans* preys on *D. discoideum* in the laboratory but does not encounter it in nature because *C. elegans* lives on rotten fruit. The nematode *Oscheius tipulae* is abundant in the soil.

**Questions:** Do the defensive bacteria that farmers carry also protect farmers from nematodes? Is this protection specific to nematodes that reside with *D. discoideum*?

**Hypotheses:** Many organisms evolve defensive mutualisms against predators. The natural habitat of *D. discoideum* is populated with nematodes. Therefore, we hypothesize that farming *D. discoideum* clones use non-food bacteria for protection from nematodes. We predicted higher fitness of farmers than non-farmers in the presence of nematodes. We also predicted to see this change of fitness only in the presence of the soil nematode, *O. tipulae*.

**Organisms:** Amoeba *Dictyostelium discoideum*, nematodes *Caenorhabditis elegans* and *Oscheius tipulae*, bacteria *Klebsiella pneumoniae* and *Burkholderia xenovorans*.

**Methods:** We compared spore production of *D. discoideum* farmers and non-farmers with and without nematodes. We also looked at nematode proliferation in the presence of farmers, non-farmers, *K. pneumoniae*, and *B. xenovorans*.

**Results:** Overall, farmer *D. discoideum* produced fewer spores than non-farmers. There was a decrease in the spore counts in the presence of nematodes for both farmers and non-farmers. There was a significant decrease in the percentage change in spore production for the farmers in the presence of soil nematodes but not laboratory nematodes. Nematode proliferation with the laboratory nematode and soil nematode did not vary in the presence of farmers, non-farmers, *K. pneumoniae* or *B. xenovorans*.

**Conclusion:** The non-food bacteria that farmers carry do not provide defence against nematodes. In fact, it was a disadvantage for farmers to carry bacteria, since the soil nematode decreased spore production for farmers compared with non-farmers. However, the differences

between the laboratory nematode and the soil nematode are marked enough to conclude that different species of nematodes respond differently to *D. discoideum* as a food source.

*Keywords:* defensive mutualisms, *Dictyostelium discoideum*, *Caenorhabditis elegans*, *Oscheius tipulae*, bacteria.

## INTRODUCTION

In symbiotic mutualisms, both partners of different species benefit from evolved interactions (Way, 1963; Bronstein, 1994; West and Herre, 1994; Hooper and Gordon, 2001; Sakai, 2002; White and Torres, 2009). For example, pea aphids (*Acyrtosiphon pisum*) provide shelter and protection to bacteria *Buchnera aphidicola*. The bacteria, in exchange, produce nutrients the aphids lack in their diet (Moran *et al.*, 2005a, 2005b). There are some cases, classified as protection mutualisms or defensive mutualisms, in which the host in the interaction is protected from predators, pathogens, or biotic and abiotic stressors by the symbiont (White and Torres, 2009).

A classic example of a defensive mutualism is *Pseudomyrmex* ants, which ward off predators of *Acacia* trees, receiving in return food and shelter (Janzen, 1966). Another classic defensive mutualism is the relationship between bacteria and fungus-growing ants (Currie *et al.*, 1999). A parasitic fungus, *Escovopsis*, sometimes threatens the ants' fungus gardens. Bacteria of the genus *Pseudonocardia* are found on the cuticle of these ants and have been shown to suppress the growth of *Escovopsis* (Currie *et al.*, 1999a, 1999b; Currie and Stuart, 2001; Poulsen *et al.*, 2009; but see Kost *et al.*, 2007; Mueller, 2012).

Another example of a bacteria–eukaryote protective mutualism is between bacteria and beetles. The bacterium, *Pseudomonas* sp., has been shown to be associated with the pederin-producing beetle, *Paederus sabaesus* (Kellner, 2001, 2002; Piel, 2002; Oliver and Moran, 2009). Piel (2002) discovered that the genes that are used to produce pederin are located on a cluster found in the genome of a *Pseudomonas* that is only found in pederin-producing beetles. Pederin is a toxin that can create painful lesions on human skin (Gelmetti and Grimalt, 1993; Oliver and Moran, 2009). Not only can pederin harm humans, but it harms beetles' predator, wolf spiders (Kellner and Dettner, 1996; Oliver and Moran, 2009).

In this study, we tested for the presence of a defensive mutualism between bacteria and the social amoeba *Dictyostelium discoideum* that can deter predators. *Dictyostelium discoideum* is a soil-dwelling microbe, which lives most of its life as single-celled amoebae that eat bacteria (Raper, 1937; Kessin, 2001). When bacteria are sparse, the social stage begins and thousands of amoebae aggregate to form a slug. The slug develops into a fruiting body in which 20% of the cells die to form a sterile stalk to hold aloft the other cells as fertile spores in a fruiting body (Kessin, 2001; Strassmann and Queller, 2010). Recently, Brock *et al.* (2011) revealed that amoebae from many clones of *D. discoideum* not only prey upon bacteria, but also farm them in a kind of primitive agriculture. The bacteria are not completely consumed by the amoebae; instead, some are kept alive through the social stage, transported to new locations with *D. discoideum* spores, and seeded out to start new prey populations (Brock *et al.*, 2011). The farming clones of *D. discoideum* carry several different food bacteria, including the one we provide as food in the laboratory, *Klebsiella pneumoniae*. They also carry some bacteria that are not good food, but instead produce effective weapons against non-farmer clones (Brock *et al.*, 2013; Stallforth *et al.*, 2013). *Dictyostelium discoideum* strains that are capable of farming bacteria thrive in environments where preferred bacteria are

scarce, but these strains do less well in laboratory conditions when bacteria are plentiful (Brock *et al.*, 2011).

Thus, some of the bacteria farmed by *D. discoideum* harm competing non-farmer clones and keep them from eating both the protecting bacteria as well as the host's edible, farmed bacteria. But it is unclear whether this is the evolved function of the symbiosis. In particular, it is not obvious how the bacteria might evolve to harm non-farming clones of *D. discoideum* without also harming their host of the same species. One possibility is that the protecting bacteria originally performed a different function. For example, they may have initially provided protection against predators.

We hypothesize that one of these non-food bacteria, *Burkholderia xenovorans*, provides protection against predators. A likely predator is a nematode, since they have been demonstrated to eat *D. discoideum* (Kessin *et al.*, 1996). Nematodes are multicellular, eukaryote soil-dwelling predators of *D. discoideum* amoebae and bacteria. We tested two nematode species, the model organism *Caenorhabditis elegans* and a wild-collected nematode, *Oscheius tipulae*, found in soil samples also containing *D. discoideum*. *Caenorhabditis elegans* has a sequenced genome, a three-day life cycle in the laboratory, and reproduction by self-fertilizing hermaphrodites. It occurs in soil and on rotting fruit (Caswell-Chen *et al.*, 2005; Félix and Braendle, 2010). It has been shown to eat *D. discoideum* (Kessin *et al.*, 1996). *Oscheius tipulae* is the most common soil nematode (Félix, 2006) and may therefore be a natural predator of *D. discoideum*.

We were interested in *B. xenovorans* because *Burkholderia* species have been shown to be toxic to nematodes (O'Quinn *et al.*, 2001). Additionally, *B. xenovorans* does not serve as food. It is found in several different *D. discoideum* farmer clones and it harms non-farmer clones (Brock *et al.*, 2013). We tested the 'bacteria protection against nematodes' hypothesis that *D. discoideum* clones have an evolved mutualism with *B. xenovorans* at least partly to provide defence against predation by nematodes. We predicted that *B. xenovorans*-carrying clones, farmers, would have higher fitness in the presence of nematodes than those *D. discoideum* clones that do not carry the bacterium, non-farmers. We measured fitness as spore counts in the presence and absence of nematodes. Additionally, we measured nematode proliferation on farmers and non-farmers. We predicted that proliferation would decrease in the presence of *D. discoideum* farmer clones. To test our hypothesis that bacteria directly harm nematodes, we measured nematode proliferation in the presence of the non-food *B. xenovorans* and the laboratory food *K. pneumoniae*. We predicted that nematode proliferation would decrease in the presence of *B. xenovorans*.

## MATERIALS AND METHODS

### Study animals

We used *C. elegans* N2 hermaphrodites donated from the Nonet lab at Washington University in St. Louis. We reared N2 strains on Nutrient Growth Media (NGM) (2.5 g bacto peptone, 3 g NaCl, 1 mL of 5 mg/mL cholesterol, 25 mL 1 M KPO<sub>4</sub> pH 6.0, 1 mL 1 M CaCl<sub>2</sub>, 1 mL 1 M MgSO<sub>4</sub> in 1 litre of ddH<sub>2</sub>O) on 60-mm diameter plates seeded with *K. pneumoniae*. We extracted *O. tipulae* isolates from plating diluted soil onto Hay agar plates (15 g of cut grass per 1.5 L, 1.5 g KH<sub>2</sub>PO<sub>4</sub>, 0.62 g Na<sub>2</sub>HPO<sub>4</sub>, and 15 g agar per litre of hay infusion) with charcoal and *K. pneumoniae*. We reared the *O. tipulae* from soil isolates on 60-mm diameter water agar plates (1% agar in water + 5 µg/mL cholesterol) (Barriere and

Félix, 2006) seeded with *K. pneumoniae*. We sequenced the internal transcribed spacer region of the ribosomal DNA of *O. tipulae* using the methods described by Barrière and Félix (2006) to identify our wild-collected nematodes. We chose L4 nematode larvae from plates with different developmental stages for both species in order to start with unfertilized young adults.






We used all of the strains of *D. discoideum* clones previously used by Brock *et al.* (2011). In our hands, only seven of the sub-cloned strains were farmers. We determined this by testing all clones for ability to carry bacteria through the social stage as spores. We placed nine individual fruiting bodies on a single SM/5 nutrient agar plate (2 g peptone, 0.2 g yeast extract, 2 g glucose, 1.9 g  $\text{KH}_2\text{PO}_4$ , 1.3 g  $\text{K}_2\text{HPO}_4$ , 0.2 g  $\text{MgSO}_4$  anhydrous, 17 g agar per litre of ddH<sub>2</sub>O) at nine different spots and waited about four days to reveal amoebae proliferation, something that could only occur if the spores carried bacteria with them. We called clones farmers if over 70% of the individual sori showed bacteria growth; we called clones non-farmers if fewer than 15% showed bacteria growth. There was only one intermediate value. In fact, of the 72 different spot tests of nine fruiting bodies each, only seven instances showed non-conforming spots: four had only a single non-conforming spot, two had two of nine non-conforming spots, and one had three of nine non-conforming spots, indicating the robustness of the test. We grew *D. discoideum* clones on SM/5 nutrient agar and used *K. pneumoniae* as their food source (100-mm diameter plates). We harvested spores after a week and used them for the experiment.

### Experimental design


We had six experimental conditions (Fig. 1). We plated nematodes with the food bacterium *K. pneumoniae* (Fig. 1, condition 1) or with the putative defensive bacterium *B. xenovorans* (Fig. 1, condition 2). Our ‘bacteria directly harms nematodes’ hypothesis predicts that *B. xenovorans* will have a negative direct effect on nematodes. Therefore, condition 1 should yield a greater number of nematodes than condition 2.


Next, we plated farmer clones carrying both non-food and their food bacterium *K. pneumoniae* without (Fig. 1, condition 3) and with nematodes (Fig. 1, condition 4). We similarly plated non-farmer clones and their food bacterium *K. pneumoniae* without (Fig. 1, condition 5) and with nematodes (Fig. 1, condition 6). Our ‘bacteria protection against nematodes’ hypothesis assumes that nematodes are predators of *D. discoideum*; therefore, conditions 3 and 4 allow us to determine the effect of nematodes on farmers, while conditions 5 and 6 allow us to determine the effect of nematodes on non-farmers. According to our ‘bacteria protection against nematodes’ hypothesis, the effect of nematodes on farmers should be more negative than the effect on non-farmers, which should be protected by their bacteria. Additionally, we compared nematode proliferation between conditions 4 and 6. According to the ‘bacteria protection against nematodes’ hypothesis, we should have a higher number of nematodes in condition 6 than in condition 4, since the nematodes should be harmed and unable to reproduce when in contact with farmers and their bacteria.


We used seven farmer clones and 17 non-farmer clones. We plated 10 L4 hermaphrodite nematodes per 23.75 cm<sup>2</sup>,  $1 \times 10^4$  *D. discoideum* spores per 23.75 cm<sup>2</sup>, and 50 µL of *K. pneumoniae* (OD<sub>600</sub> = 1.5) or *B. xenovorans* (OD<sub>600</sub> = 1.5) depending on the treatment or control. For example, for the first treatment we plated 10 L4 hermaphrodites per 23.75 cm<sup>2</sup> and 50 µL of *B. xenovorans* (OD<sub>600</sub> = 1.5). SM/5 nutrient agar plates were used for the *C. elegans* experiment. Hay agar plates were used for the *O. tipulae* experiment. We could


Conditions	<i>K. pneumoniae</i>	<i>B. xenovorans</i>	Farmers	Non-Farmers	Nematodes
 1	✓				✓
 2		✓			✓
 3	✓		✓		
 4	✓		✓		✓
 5	✓			✓	


Legend

*K. pneumoniae* 

*B. xenovorans* 

Farmers 

Non-Farmers 

Nematodes 

**Fig. 1.** Design of the experiment with complete list of controls and treatments. We had two plates for each control and treatment, which were performed simultaneously for a total of 40 plates per block. We performed each block twice.

not use the same type of agar plate for both species of nematodes because *O. tipulae* did not grow well on SM/5 plates presumably because of the overwhelming amount of nutrients compared with wild conditions. Therefore, we used the more nutrient poor hay agar plates, which more closely mimic the nutrients found in the soil.

We collected data on nematode number and *D. discoideum* spore production after 5 days to restrict our results to the first generation of nematodes. For each *D. discoideum* clone and nematode combination, all six treatments were performed twice. These were done in blocks of eight *D. discoideum* clones at a time. After chilling the plates on ice for about 10 minutes to stop nematode movement and ensure accurate counting, we counted nematodes using a Nikon SMZ 1500 dissecting scope. We placed a transparent grid (2.54 × 2.54 cm) with 10 random marked squares (0.506 × 0.506 cm) underneath the petri plate and counted nematodes only found on the marked squares. We harvested *D. discoideum* spores from plates using 7 mL of KK2 and a sterilized spatula and counted spores using a hemocytometer.

To visualize nematode feeding, we fed nematodes for about 4 hours on *K. pneumoniae* transformed to express green fluorescent protein (GFP) and *D. discoideum* spores and cells transformed to express red fluorescent protein (RFP). Afterwards, we paralysed the nematodes with 10 mM Na-azide and placed the nematodes on slides. We took pictures on a Nikon Eclipse E1000 with RFP and GFP filters.

### Statistical analyses

#### *D. discoideum* spore counts with and without nematodes


For all statistical testing, we used the R statistical computing package v.2.14.1 (R Development Core Team, 2012). To investigate the effect of nematodes on *D. discoideum*'s fitness, we used spore counts as our response variable. To measure the change of spore counts in the presence of nematodes, we used percentage change in spore number  $((\text{presence} - \text{absence}) / \text{absence}) \times 100$ . We used the Mann-Whitney-Wilcoxon Test to determine whether there was a difference between the farmers and non-farmers.

#### Nematode counts

To investigate the direct effect of *B. xenovorans* and *D. discoideum* on the fitness of nematodes, we looked for a significant difference between nematode counts in the presence of farmers and non-farmers, *B. xenovorans* and *K. pneumoniae*. The *C. elegans* and *O. tipulae* nematode counts were square root transformed to satisfy the assumptions for using a parametric statistical test. We used an ANOVA to test for differences in nematode counts between nematodes fed on *B. xenovorans* versus *K. pneumoniae* and also farmers versus non-farmers.

## RESULTS

We observed active consumption of *K. pneumoniae* and *D. discoideum* cells by *C. elegans* (Fig. 2), which confirms our assumption for our 'bacteria protection against nematodes' hypothesis. We observed the presence of labelled *K. pneumoniae* (Fig. 2A) and both spores and cells of *D. discoideum* (Fig. 2B, 2C) in the guts of *C. elegans*. However, we saw that *D. discoideum* spores remained intact in the nematode gut, validating the result of Kessin *et al.* (1996). Therefore, nematodes are predators of *D. discoideum* cells, and potential dispersers of *D. discoideum* spores. We also observed the behaviour of the nematodes. In the early stages of the nematodes' life cycle, L1, L2, and L2 dauers can climb the stalks of



**Fig. 2.** *C. elegans* digestion after 3 hours. (A) *C. elegans* fed on GFP-labelled *K. pneumoniae*. (B) *C. elegans* fed on RFP-labelled *D. discoideum* spores. (C) *C. elegans* fed on RFP-labelled *D. discoideum* cells.

the fruiting bodies and freely move around inside the sorus. This behaviour was also witnessed and documented in Kessin (2001).

#### *D. discoideum* spore counts with and without nematodes

To directly test our ‘bacteria protection against nematodes’ hypothesis, we compared farmers and non-farmers for the percentage change of spore counts in the presence of nematodes. Our hypothesis predicts that the change will be negative (nematodes harm spore production) but that it will be less negative for farmers because of protection by their bacteria. The presence of either *C. elegans* or *O. tipulae* nematodes produced similar results. Counter to our prediction, farmers were in fact more harmed by nematodes, significantly for one species and nearly significant and in the same direction for the other (*C. elegans*,  $P = 0.06$ , Fig. 3A; *O. tipulae*,  $P < 0.01$ , Fig. 3B).

#### Nematode counts


We also wanted to test our ‘bacteria directly harms nematodes’ hypothesis by quantifying whether there is a direct negative effect of non-food bacterium, *B. xenovorans* on *D. discoideum* predators, particularly nematodes. If *B. xenovorans* has a direct negative effect on nematodes, we expected the proliferation of nematodes to be lower when nematodes are fed strictly on *B. xenovorans* than on *K. pneumoniae*. For our ‘bacteria protection against nematodes’ hypothesis not to be rejected, the proliferation of nematodes should be lower in the presence of farmers than non-farmers, since farmers carry *B. xenovorans* that could potentially harm nematodes. *Caenorhabditis elegans* proliferation was consistent between the treatments (ANOVA: *K. pneumoniae* vs. *B. xenovorans*,  $P = 0.898$ ; farmers vs. non-farmers,  $P = 0.678$ ; Fig. 4A). Similarly, *O. tipulae* proliferation was consistent between treatments (ANOVA: *K. pneumoniae* vs. *B. xenovorans*,  $P = 0.223$ ; farmers vs. non-farmers,  $P = 0.09$ ; Fig. 4B). Although the farmer versus non-farmer comparison approached significance, it was in the opposite direction to our prediction, with nematodes faring better on farmers.

## DISCUSSION

A previous study showed that farmer clones of *D. discoideum* are able to thrive in environments where non-farmers would do poorly, particularly when good food bacteria are scarce (Brock *et al.*, 2011). Therefore, there is a positive effect of carrying the bacteria through the social stage. We know that the bacteria are able to help in host conspecific competition and that they do so by producing compounds that harm competing non-farmer clones (Brock *et al.*, 2013; Stallforth *et al.*, 2013). Here we tested a related hypothesis. If *B. xenovorans* produces compounds that are harmful to non-farming *D. discoideum* clones, they might also serve as a weapon to fight against predatory nematodes. If they protect against nematodes, then the protection against non-farmer clones might be a side-effect rather than an adaptation. However, the predator-protection hypothesis was not supported.


In order to measure *D. discoideum* fitness, we looked at percentage change of spore production as a measure of fecundity in the presence versus absence of predators. Both nematodes, *C. elegans* and *O. tipulae*, harmed fecundity for both farmers and non-farmers. Under the ‘bacteria protection against nematodes’ hypothesis, farmers should decrease less than non-farmers in the presence of nematodes. We observed similar results when we used different species of nematodes. With *C. elegans*, there was no significant percentage change





**Fig. 3.** Mean percentage change ( $\pm$ s.d.) in spore production of non-farmer ( $n = 17$ ) and farmer ( $n = 7$ ) *D. discoideum* clones in the presence of nematode predators: (A) *C. elegans* (Mann-Whitney-Wilcoxon Test, non-farmer vs. farmer:  $W = 184$ ,  $P > 0.06$ ); (B) *O. tipulae* (Mann-Whitney-Wilcoxon Test, non-farmer vs. farmer:  $W = 134$ ,  $P < 0.01$ ).

for farmers, which rejects our hypothesis. Furthermore, with *O. tipulae*, there was a significant percentage change of spore counts for farmers; however, the decrease was much higher than for non-farmers, which also rejects our hypothesis. This allows us to conclude that the farmers' bacteria do not help protect the farmers against nematodes. On the contrary, the bacteria tend to increase the farmers' risk of predation. One hypothesis is that it is due to the bacteria being harboured in the sorus. They may therefore attract the nematodes



**Fig. 4.** Nematode counts ( $\pm$ S.D.) of (A) *C. elegans* and (B) *O. tipulae* (farmers,  $n = 7$ ; non-farmers,  $n = 17$ ; *B. xenovorans*,  $n = 2$ ; *K. pneumoniae*,  $n = 6$ ). We replicated and duplicated each treatment and control. We performed an ANOVA. For *C. elegans*: *K. pneumoniae* vs. *B. xenovorans*,  $P = 0.898$ ; non-farmers vs. farmers,  $P = 0.678$ . For *O. tipulae*: *K. pneumoniae* vs. *B. xenovorans*,  $P = 0.223$ ; non-farmers vs. farmers,  $P = 0.09$ .

to this structure where they consume spores in addition to the bacteria. To further test our 'bacteria protection against nematodes' hypothesis, we measured proliferation of nematodes on farmers and non-farmers. Neither *C. elegans* nor *O. tipulae* proliferation changed when reared on *D. discoideum* farmers or *D. discoideum* non-farmers. Therefore, the non-food bacteria do not protect *D. discoideum* farmers from predation.

As for our ‘bacteria directly harms nematodes’ hypothesis, we found that for both species of nematodes, *C. elegans* and *O. tipulae*, proliferation did not change when reared on *K. pneumoniae* versus in the presence of non-food bacteria, *B. xenovorans*. We can conclude that *B. xenovorans* does not harm the proliferation of *C. elegans* or *O. tipulae*, contrary to our prediction (Fig. 4). The non-food bacteria do not directly harm either nematode, *C. elegans* or *O. tipulae*, which contrasts with the finding that *C. elegans* was killed by *Burkholderia psuedomallei* (O’Quinn *et al.*, 2001).

In conclusion, all predictions based on the hypothesis that carrying *B. xenovorans* protects farmers against nematodes failed. Therefore, the bacteria carried by farmers are not used for protection against *C. elegans* and *O. tipulae*. This lack of an effect against predators makes it more likely that the true function of carrying *B. xenovarans* is the previously demonstrated effect in aiding the farmers’ competition between conspecifics (Brock *et al.*, 2013; Stallforth *et al.*, 2013).

### ACKNOWLEDGEMENTS

We thank Scott Dour and the Nonet lab for providing us with the N2 *C. elegans* strain and for helping us with culturing techniques. Wild collections of nematodes were made by Debbie Brock, Ben Howard, and Emily McMeekin. We would also like to thank Debbie Brock for providing us with the clones of *D. discoideum*, and the entire Strassmann-Queller Laboratory for their helpful comments and insights. This material is based on work supported by the John Templeton Foundation grant no. 43677, NSF grant nos. DEB 1011513, DEB 1146375, and IOS 1256416.

### REFERENCES

- Barriere, A. and Félix, M.-A. 2006. Isolation of *C. elegans* and related nematodes. *WormBook: The Online Review of C. elegans Biology* [<http://www.ncbi.nlm.nih.gov/books/NBK19764/>].
- Brock, D.A., Dougals, T.E., Queller, D.C. and Strassmann, J.E. 2011. Primitive agriculture in a social amoeba. *Nature*, **469**: 393–396.
- Brock, D.A., Read, S., Bozhchenko, A., Queller, D.C. and Strassmann, J.E. 2013. Social amoeba farmers carry defensive symbionts to protect and privatize their crops. *Nature Commun.*, **4**: 1–7.
- Bronstein, J.L. 1994. Our current understanding of mutualism. *Q. Rev. Biol.*, **69**: 31–51.
- Caswell-Chen, E.P., Chen, J., Lewis, E.E., Douhan, G.W., Nadler, S.A. and Carey, J.R. 2005. Revising the standard wisdom of *C. elegans* natural history: ecology of longevity. *Sci. Aging Knowledge Environ.*, **40**: pe30.
- Currie, C.R. and Stuart, A.E. 2001. Weeding and grooming of pathogens in agriculture by ants. *Proc. R. Soc. Lond. B*, **268**: 1033–1039.
- Currie, C.R., Mueller, U.G. and Malloch, D. 1999a. The agricultural pathology of ant fungus gardens. *Proc. Natl. Acad. Sci. USA*, **96**: 7998–8002.
- Currie, C.R., Scott, J.A., Summerbell, R.C. and Malloch, D. 1999b. Fungus-growing ants use antibiotic-producing bacteria to control garden parasites. *Nature*, **398**: 701–704.
- Félix, M.-A. 2006. *Oscheius tipulae*. *WormBook: The Online Review of C. elegans Biology* [[http://www.wormbook.org/chapters/www\\_genomesOtipulae/genomesOtipulae.html](http://www.wormbook.org/chapters/www_genomesOtipulae/genomesOtipulae.html)].
- Félix, M.-A. and Braendle, C. 2010. The natural history of *Caenorhabditis elegans*. *Curr. Biol.*, **20**: R965–R969.
- Gelmetti, C. and Grimalt, R. 1993. Paederus dermatitis: an easy diagnosable but misdiagnosed eruption. *Eur. J. Pediatrics*, **152**: 6–8.
- Hooper, L.V. and Gordon, J.I. 2001. Commensal host–bacterial relationships in the gut. *Science*, **292**: 1115–1118.

- Janzen, D.H. 1966. Coevolution of mutualism between ants and acacias in Central America. *Evolution*, **20**: 249–275.
- Kellner, R.L.L. 2001. Horizontal transmission of biosynthetic capabilities for pederin in *Paederus melanurus* (Coleoptera: Staphylinidae). *Chemoecology*, **11**: 127–130.
- Kellner, R.L.L. 2002. Molecular identification of an endosymbiotic bacterium associated with pederin biosynthesis in *Paederus sabaues* (Coleoptera: Staphylinidae). *Insect Biochem. Mol. Biol.*, **32**: 389–395.
- Kellner, R.L.L. and Dettner, K. 1996. Differential efficacy of toxic pederin in deterring potential arthropod predators of *Paederus* (Coleoptera: Staphylinidae) offspring. *Oecologia*, **107**: 293–300.
- Kessin, R.H. 2001. *Dictyostelium: Evolution, Cell Biology, and Development of Multicellularity*. Cambridge: Cambridge University Press.
- Kessin, R.H., Gundersen, G.G., Zaydfudium, V. and Grimson, M. 1996. How cellular slime molds evade nematodes. *Proc. Natl. Acad. Sci. USA*, **93**: 4857–4861.
- Kost, C., Lakatos, T., Böttcher, I., Arendholz, R., Redenbach, M. and Wirth, R. 2007. Non-specific association between filamentous bacteria and fungus-growing ants. *Naturwissenschaften*, **94**: 821–828.
- Moran, N.A., Degnan, P.H., Santos, S.R., Dunbar, H.E. and Ochman, H. 2005a. The players in a mutualistic symbiosis: insects, bacteria, viruses, and virulence genes. *Proc. Natl. Acad. Sci. USA*, **102**: 16919–16926.
- Moran, N.A., Tran, P. and Gerardo, N.M. 2005b. Symbiosis and insect diversification: an ancient symbiont of sap-feeding insects from the bacterial phylum Bacteroidetes. *Appl. Environ. Microbiol.*, **71**: 8802–8810.
- Mueller, U.G. 2012. Symbiont recruitment versus ant-symbiont co-evolution in the attine ant-microbe symbiosis. *Curr. Opin. Microbiol.*, **15**: 269–277.
- Oliver, K. and Moran, N. 2009. Defensive symbionts in aphids and other insects. In *Defensive Mutualism in Microbial Symbiosis* (J.F. White, Jr. and M.S. Torres, eds.), pp. 129–148. Boca Raton, FL: CRC Press.
- O’Quinn, A.L., Wiegand, E.M. and Jeddelloh, J.A. 2001. *Burkholderia pseudomallei* kills the nematode *Caenorhabditis elegans* using an endotoxin-mediated paralysis. *Cell. Biol.*, **6**: 381–393.
- Piel, J. 2002. A polyketide synthase-peptide synthetase gene cluster from an uncultured bacterial symbiont of *Paederus* beetles. *Proc. Natl. Acad. Sci. USA*, **99**: 14002–14007.
- Poulsen, M., Little, A. and Currie, C. 2009. Fungus-growing ant-microbe symbiosis. In *Defensive Mutualism in Microbial Symbiosis* (J.F. White, Jr. and M.S. Torres, eds.), pp. 149–166. Boca Raton, FL: CRC Press.
- Raper, K.B. 1937. Growth and development of *Dictyostelium discoideum* with different bacterial associates. *J. Agric. Res.*, **55**: 289–316.
- R Development Core Team. 2012. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Sakai, S. 2002. A review of brood-site pollination mutualism: plants providing breeding sites for their pollinators. *J. Plant Res.*, **115**: 161–168.
- Stallforth, P., Brock, D.A., Cantley, A.M., Tian, X., Queller, D.C., Strassmann, J.E. et al. 2013. A bacterial symbiont is converted from an inedible producer of beneficial molecules into food by a single mutation in the *gacA* gene. *Proc. Natl. Acad. Sci. USA*, **110**: 14528–14533.
- Strassmann, J.E. and Queller, D.C. 2010. The social organism: congresses, parties, and committees. *Evolution*, **64**: 605–616.
- Way, M.J. 1963. Mutualism between ants and honeydew-producing Homoptera. *Annu. Rev. Entomol.*, **8**: 307–344.
- West, S.A. and Herre, E.A. 1994. The ecology of the New World fig-parasitizing wasps *Idarnes* and implications for the evolution of the fig-pollinator mutualism. *Proc. R. Soc. Lond. B*, **258**: 67–72.
- White, J. and Torres, M. 2009. Introduction: symbiosis, defensive mutualism, and variations on the theme. In *Defensive Mutualism in Microbial Symbiosis* (J.F. White, Jr. and M.S. Torres, eds.), pp. 3–8. Boca Raton, FL: CRC Press.

