

2011

Will the use of Less Fecund Cultivars Reduce the Invasiveness of Perennial Plants?

Tiffany M. Knight

Washington University in St Louis, tknight@wustl.edu

Kayri Havens

Chicago Botanic Garden

Pati Vitt

Chicago Botanic Garden

Follow this and additional works at: http://openscholarship.wustl.edu/bio_facpubs



Part of the [Biology Commons](#)

Recommended Citation

Knight, Tiffany M.; Havens, Kayri; and Vitt, Pati, "Will the use of Less Fecund Cultivars Reduce the Invasiveness of Perennial Plants?" (2011). *Biology Faculty Publications & Presentations*. Paper 17.

http://openscholarship.wustl.edu/bio_facpubs/17

This Article is brought to you for free and open access by the Biology at Washington University Open Scholarship. It has been accepted for inclusion in Biology Faculty Publications & Presentations by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

Will the use of Less Fecund Cultivars Reduce the Invasiveness of Perennial Plants?

Author(s): Tiffany M. Knight, Kayri Havens and Pati Vitt

Source: BioScience, 61(10):816-822. 2011.

Published By: American Institute of Biological Sciences

URL: <http://www.bioone.org/doi/full/10.1525/bio.2011.61.10.11>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Will the Use of Less Fecund Cultivars Reduce the Invasiveness of Perennial Plants?

TIFFANY M. KNIGHT, KAYRI HAVENS, AND PATI VITT

*Many invasive species were originally introduced for horticultural purposes, and several continue to be profitable for the green (nursery, horticulture, and landscape) industry. Recently, some plant suppliers have marketed less fecund cultivars of several invasive species, including glossy buckthorn (*Frangula alnus*), burning bush (*Euonymus alatus*), and Japanese barberry (*Berberis thunbergii*), as “safe” alternatives to invasive relatives. We use published matrix population models to simulate the effect of reducing fecundity on the population growth rates of invasive species. We show that large changes in fecundity result in relatively small changes to the population growth rates of long-lived species, which suggests that less fecund cultivars may still provide an invasive threat. Furthermore, many cultivars are clonal selections, and if crossed with other cultivars or selfed, they produce offspring with traits and fecundities that do not resemble the parent plant. On the basis of these two lines of evidence, we suggest that only female sterile cultivars that cannot reproduce asexually should be considered “safe” and noninvasive. Marketing less fecund cultivars as “safe” is premature at this time, and further research is necessary to determine the potential invasiveness of different cultivars.*

Keywords: invasive plant, ornamental plant, cultivar, demography

Invasive woody plants pose serious threats to native species and ecosystems (Wilcove et al. 1998), and 82% of the invasive trees and shrubs in the United States were introduced through horticulture (Reichard 1997, Reichard and White 2001). The high proportion of invasive species originating from horticulture is not surprising, considering that many of the traits that we desire in ornamental plants are the same traits that are found in invasive species; for example, they are likely to form dense stands (i.e., are good for erosion control), are adaptable to a wide range of landscape conditions, have high levels of flower production and a long flowering period (i.e., they are attractive in gardens), have a high fruit set (especially of fleshy fruit, which can be attractive to bird and animal dispersers), and are able to spread on their own vegetatively or by seed (Kitajima et al. 2006).

Once a species is identified as invasive to natural areas, it is important to prevent the sale, and thus the spread, of that species. However, this effort is complicated by the high demand for the species by consumers and the high economic value of the species to the green (nursery, horticulture, and landscape) industry. For example, Japanese barberry (*Berberis thunbergii*) is considered invasive in 30 eastern and central states in the United States, and many states are considering banning, or have already banned, the sale of this species (Harrington et al. 2003). The loss of Japanese barberry is vexing to many in the green industry, since this species is a valuable commodity. In the state of

Connecticut alone, annual sales of the shrub are estimated to be \$5 million (Lehrer et al. 2006).

Species sold in the green industry often have several cultivars that may vary dramatically in their seed production and viability. In most cases, these cultivars were created to provide options for landscape designs and not to create less invasive forms of the species. For example, cultivars selected to produce more petals often produce fewer seeds, and hybrid cultivars formed by crossing distantly related taxa can produce fewer seeds or can be sterile (Ramney 2004). It is only recently that many researchers are purposely creating cultivars of invasive plants that are less fecund, in an effort to reduce their invasive potential (e.g., Ranney 2004, Drew et al. 2010). Regardless of how and why the cultivar was formed, if it is unlikely to invade and harm natural areas, these cultivars should be exempt from invasive species bans. The critical question then becomes how much of a reduction in seed production or seed viability is necessary to create a cultivar that will not be invasive in natural areas. The answer depends largely on the demography of the plant across its entire life cycle, as well as the sensitivity of its population growth rate to reductions in seed production. In a worst-case scenario, large reductions in the seed production of a cultivar relative to that of the wild type would result in relatively small changes in the population growth rate of the species in the wild and therefore to the invasiveness of the plant. To date, no studies have quantified the demography of

different cultivars across their life cycle and compared their projected population growth rates.

Offspring of a cultivar might perform much differently from its horticulturally produced parent plants. For taxa that do not spread vegetatively, cultivars per se do not invade: Their offspring invade. Cultivar offspring may or may not resemble the parents, because cultivars are created from sports (somatic mutations that are selected and propagated clonally) or may be clones of complex, multispecies or multicultivar hybrids. In either case, the offspring will not “breed true.” Very few ornamental taxa are “seed selections” of inbred lines that will produce morphologically similar offspring when selfed or crossed with another individual of the same cultivar. Cultivars of long-lived perennial plants are not created by seed selections because the time required makes this method impractical. Furthermore, there is a financial incentive to produce clonal cultivars rather than seed selections because the clonal cultivars will not breed true, which would give the breeder built-in property rights and would secure monetary returns (Drew et al. 2010). In addition, in garden settings, there may be more than one cultivar of a species present or there may be a nearby feral population of the species. In both cases, crossing between cultivars or between a cultivar and an invasive wild type could produce offspring dissimilar from one or both parents.

We currently do not have demographic studies across the entire life cycle for species that are economically important to the green industry (e.g., Japanese barberry), and therefore, we cannot quantify how much of a reduction in seed production or seed viability is necessary to create a cultivar incapable of invading natural areas. However, we can synthesize demographic studies on wild invasive plant species as a first step in considering the degree to which reductions in seed production might result in reductions in population growth rates. In our review of the literature, we found that matrix population models have been constructed for 24 invasive plant species; these studies quantify the demography of wild invasive plant species in natural areas of their invasive ranges.

In the present article, we will first illustrate that cultivars of invasive plants are being advertised as noninvasive on the basis of limited scientific evidence. Second, we will quantitatively synthesize matrix population models parameterized for wild invasive plants and will evaluate how the population growth rates of these plants would change under a range of reductions in seed production or seed viability. Third, we will discuss the research needed to evaluate whether a cultivar is likely to be safe to natural areas. We suggest that marketing of cultivars as “safe” or “noninvasive” needs to be based on evaluations that consider the entire life cycle of the cultivar and its offspring.

Examples of three invasive species with “noninvasive” cultivars

Here, we highlight three species that (a) are popular with consumers and are therefore highly profitable for the green

industry, (b) have wild types that are natural-area invaders, and (c) have cultivars with dramatically lower rates of seed production or seed viability. For each species, there is experimental evidence that some cultivars have lower fecundity than the invasive wild type. Specifically, *Frangula alnus* (‘Fine Line’ glossy buckthorn), *Euonymus alatus* (‘Rudy Haag’ burning bush), and *B. thunbergii* (‘Crimson Pygmy’ Japanese barberry) are cultivars with lower rates of seed production or seed viability than many available on the market. Recently, some plant suppliers have begun to market these cultivars as noninvasive. As an illustration of this, we provide examples from Web sites and other sources. We discuss the reasons for which marketing cultivars such as these as safe to natural areas is premature.

Buckthorn. *Frangula alnus* (formerly *Rhamnus frangula*; Rhamnaceae), which is originally from Europe, forms dense thickets in North American forest understories and reduces native herbaceous plant abundance and inhibits tree regeneration (Fagan and Peart 2004, Frappier et al. 2004). ‘Fine Line’ buckthorn originated from cross-pollination of a female ‘Columnaris’ cultivar and a male ‘Asplenifolia’ cultivar (Williams 2004); individuals have the feathery foliage of ‘Asplenifolia’ with the narrow upright habit of ‘Columnaris.’ In an experimental study, seeds of ‘Fine Line’ had significantly lower germination than the wild-type buckthorn (12% and 83.7%, respectively; Mark C. Starrett, Department of Plant and Soil Sciences, University of Vermont, Burlington, personal communication, 28 December 2010). Propagation of the ‘Fine Line’ cultivar for sale has been achieved entirely through asexual reproduction by cuttings (Williams 2004). It is unknown whether the offspring produced by seed by ‘Fine Line’ will have a morphology or a demographic performance similar to those of the parent plant. However, since this cultivar is of hybrid origin, it is likely that the offspring will not breed true.

‘Fine Line’ buckthorn is advertised as a noninvasive by several Web sites. For example, this cultivar was featured as an “Indiana Plant Pick” in the November–December 2009 issue of the Indiana Nursery and Landscapers Association Journal (see Jacquart and Knight 2010 for a response to this choice). Furthermore, one Web site claims that “It has been proven to be an environmentally friendly plant that is not invasive like other types of buckthorn” (7 April 2010; www.colorchoiceplants.com/fine_line.htm#ixzz0htbHfEgz).

Burning bush. *Euonymus alatus* (Celastraceae), which is native to East Asia, forms dense thickets and displaces native plant species in forest understories of the United States (Marinelli and Hanson 1996). The ‘Rudy Haag’ cultivar is a more compact form of the popular ‘Compactus’ cultivar. The origin of ‘Rudy Haag’ has not been documented, but it likely resulted from a seedling or a sport from ‘Compactus’ (Mark Brand, Department of Plant Science and Landscape Architecture, University of Connecticut, Storrs, personal communication, 24 May 2011).

Recent studies show lower seed and fruit production in 'Rudy Haag' than in the 'Compactus' cultivar. In field plots, 'Rudy Haag' produced an average of 12 seeds per plant compared with an average of 1238 produced by the 'Compactus' cultivar (Cynthia Finneseth, College of Agriculture, University of Kentucky, Lexington, personal communication, 31 March 2011). In another study, 12% of the flowers produced fruit in 'Compactus' compared with 0.5% in 'Rudy Haag' (Ingram et al. 2004). Taken together, the 'Rudy Haag' cultivar yields a 96%–99% decrease in reproduction relative to the 'Compactus' cultivar. However, when 'Rudy Haag' is grown with other cultivars, it can cross-pollinate with them and have relatively high fruit production (Mark Brand, Department of Plant Science and Landscape Architecture, University of Connecticut, Storrs, personal communication, 24 May 2011).

The 'Rudy Haag' cultivar has been called less invasive by some. It was picked as the plant of the month by Green Sense for being a "noninvasive burning bush" (<http://greensenseassess.com/2010/09/07/september-plant-of-the-month>). In a recent newspaper article, a horticulturist for the Washington State University Benton County Extension Office stated that "with very limited seed production, 'Rudy Haag' is much less of an invasive threat" (Ophardt 2009).

Japanese barberry. *Berberis thunbergii* (Berberidaceae), which is native to Japan, invades North American forest understories, displaces native shrubs (Kourtev et al. 1998), and causes changes in soil properties (Ehrenfeld et al. 2001). The 'Crimson Pygmy' cultivar is known for its small stature and burgundy foliage. A recent study by Lehrer and colleagues (2006) demonstrates that this cultivar produced fewer fruits (75 versus 1140 fruits per plant) and fewer seeds per plant (90 versus 1135) than the wild type.

'Crimson Pygmy' was introduced by Van Eyck in Boskoop, in the Netherlands, in 1942, likely as a seedling from *B. thunbergii* var. *atropurpurea*. Recent research indicates that 'Crimson Pygmy' is not a hybrid (Lubell et al. 2008). However, gardeners have noted that 'Crimson Pygmy' plants will not breed true from seed, and some offspring fail to have the burgundy foliage or the small size observed in the parent (<http://davesgarden.com/guides/pf/go/59576>). Furthermore, open-pollinated offspring of purple- and yellow-leaved cultivars produce at least 20% green-leaved (resembling the wild type) offspring (Lehrer et al. 2006), which indicates that these cultivars also do not breed true.

It is difficult to find scientific data on the performance of many cultivars, and some of the information in the case studies provided here are unpublished or anecdotal. In some cases, there is excellent experimental research on seed production of different cultivars or on the viability and germination rate of the seeds. However, this research is limited in its scope in that only a few demographic vital rates were considered and only horticulturally produced plants (not their offspring) were the focus. Information about other important demographic vital rates that influence the

potential growth rate and invasiveness of a species, such as the time to reproduction, is nonexistent. These case studies demonstrate a problem: Cultivars with reduced fertility are being marketed by some suppliers as less invasive, but scientific evidence for this claim is lacking.

The effects of reducing fecundity on population dynamics of invasive plants

Invasive species typically show rapid rates of population growth (Ramula et al. 2008). If cultivars can be created that have low rates of population growth in natural areas, it might be reasonable to conclude that these are less of an invasive threat. Matrix population models synthesize vital rates across the entire life cycle of a population and can be used to project the change in population size over time. The simplest form of these models assumes density independence and constant environments, and therefore, these models are not thought to be good at forecasting (Crone et al. 2011). However, the models are thought to be more useful as a comparative tool; in this context, they provide a way to synthesize vital rates and to compare the performance of different cultivars growing in different habitats or geographic regions.

Here, in order to simulate the potential effectiveness of reduced fecundity as a means to control the invasiveness of a horticultural cultivar, we found studies in which matrix population models were parameterized for invasive plant populations in their introduced range and in which how reducing fecundity (in practice, this could be achieved by reducing seed production, seed viability, or germination rates) influences the population growth rate was quantified.

To find studies, we conducted a literature search using the Web of Science (http://thomsonreuters.com/products_services/science/science_products/a-z/web_of_science) and the keyphrase *demograph** and *plant** and *matri** and *invasive*. We identified 24 species from 14 different plant families that ranged in life history from annual herbs to long-lived trees. Of these, we considered a subset of 19 species from 13 different plant families that ranged in population growth rates, λ , from 1.0 to 2.5 (table 1). Values of λ indicate whether populations are declining ($\lambda < 1$), static ($\lambda = 1$), or growing ($\lambda > 1$) through time. We used species with similar growth rates so that we could compare groups of species with different life histories without confounding the differences in population growth rates across these groups. Most studies were parameterized with a single population and year. We separated species into three categories based on their life histories: monocarpic herbs and grasses (eight species), perennial herbs and grasses (two species), and perennial shrubs and trees (nine species).

In each study, the plants were separated into discrete stage or size classes and a deterministic matrix population model was parameterized. When multiple sites or years were considered, we averaged the vital rates to create a single demographic matrix, A , for each invasive species. The matrix population model is $n_{t+1} = A \times n_t$, where the vector n_t gives the number of individuals in each stage at time t

Table 1. Invasive plant species in which matrix population models have been parameterized.

Life form category	Species	Plant family	Population growth rate (λ)	Reduction in fecundity for $\lambda \leq 1$ (percentage)	Location of study	Citation
Monocarpic	<i>Alliaria petiolata</i>	Brassicaceae	1.42	55–60	Compiled from multiple sites in the United States	Davis et al. 2006
	<i>Carduus nutans</i>	Asteraceae	1.80	60–65	New Zealand	Shea and Kelly 1998
	<i>Cirsium vulgare</i>	Asteraceae	1.54	65–70	Nebraska	Tenhumberg et al. 2008
	<i>Dipsacus sylvestris</i>	Dipsacaceae	2.33	90–95	Michigan	Werner and Caswell 1977
	<i>Echium vulgare</i>	Boraginaceae	2.28	75–80	Canberra, Australia	Grigulis et al. 2001
	<i>Heracleum mantegazzianum</i>	Apiaceae	1.21	55–60	Czech Republic	Nehrbass et al. 2006
	<i>Polygonum perfoliatum</i>	Polygonaceae	1.56	60–65	Pennsylvania	Hyatt and Araki 2006
	<i>Setaria faberi</i>	Poaceae	1.51	30–35	Iowa	Davis et al. 2004
Perennial herb or grass	<i>Centaurea maculosa</i>	Asteraceae	1.33	75–80	Michigan	Emery and Gross 2005
	<i>Molinia caerulea</i>	Poaceae	1.86	95–100	Belgium	Jacquemyn et al. 2005
Shrub or tree	<i>Ardisia elliptica</i>	Myrsinaceae	1.16	95–100	Florida	Koop and Horvitz 2005
	<i>Clidemia hirta</i>	Melastomataceae	1.78	95–100	Hawaii	DeWalt 2006
	<i>Cytisus scoparius</i>	Fabaceae	1.40	95–100	Washington state	Parker 2000
	<i>Gleditsia triacanthos</i>	Fabaceae	2.03	95–100	Cordoba Province, Argentina	Marco and Páez 2000
	<i>Parkinsonia aculeata</i>	Fabaceae	1.53	95–100	Central Australia	Raghu et al. 2006
	<i>Pinus nigra</i>	Pinaceae	1.96	95–100	New Zealand	Buckley et al. 2005
	<i>Rubus armeniacus</i>	Rosaceae	1.15	n/a	Oregon	Lambrecht-McDowell and Radosevich 2005
	<i>Syzygium jambos</i>	Myrtaceae	1.16	95–100	Puerto Rico	Brown et al. 2008

Note: λ was based on the average matrix for studies in which multiple populations and years were considered. n/a, not applicable.

(Caswell 2001). The asymptotic population growth rate, λ , is the dominant eigenvalue of A . In order to quantify how reductions in fecundity would affect λ , we multiplied all fecundity matrix elements by $(1 - p)$, where p is the proportion by which fecundity is reduced. We reduced fecundity in increments of .05 across a range of 0 (no reduction) to 1 (completely sterile).

We conducted an elasticity analysis of each unaltered matrix ($p = 0$) to determine the proportional sensitivity of λ to changes in the matrix element a_{ij} :

$$e_{ij} = \frac{a_{ij}}{\lambda} \frac{\partial \lambda}{\partial a_{ij}} = \frac{\partial(\log \lambda)}{\partial(\log a_{ij})}$$

For each species, we summed the elasticity values across all matrix elements associated with fecundity and used an ANOVA (analysis of variance) and Tukey's HSD (honestly significant difference) test to determine whether plants in each of the three life history categories described above differ in the sensitivity of λ to changes in fecundity.

This quantitative synthesis demonstrates that there are generalities in the relationship between fecundity and population growth rate attributable to plant life history. The population growth rate drops dramatically with reductions in fecundity for short-lived plant species (e.g., monocarpic species) but not for longer-lived species of herbs, shrubs, and trees (figure 1). It typically takes reductions in fecundity

between 95%–100% to reduce the population growth rates of shrubs and trees to 1 (figure 1, table 1). The population growth rates of shorter-lived species are more sensitive to changes in fecundity than those of longer-lived species (figure 2; $F(2,19) = 12.022$, $p < .001$). This result is not surprising, and several plant demography reviews have pointed out the negative relationship between life span and the sensitivity of the population growth rate to changes in fecundity (e.g., Silvertown et al. 1993), including a review focused on invasive plant species (Ramula et al. 2008). Our review highlights that reductions in seed production or seed viability alone are likely not sufficient to create a "safe" cultivar for an invasive long-lived perennial or woody plant (e.g., *F. alnus*, *E. alatus*, *B. thunbergii*) and also highlights the importance of considering the entire plant life cycle in evaluations of demographic performance and invasive potential. Our goal in these analyses was not to identify an exact magnitude by which reductions in seed production can lead to a safe cultivar but, rather, to make the point that it is difficult to decrease the population growth rate of long-lived species by reducing seed production and that one should not assume that dramatic reductions in seed production will result in similarly dramatic reductions in population growth.

In the studies reviewed here, density-independent matrix population models were employed, and thus population growth in the absence of density-dependent regulation was projected. However, invasive plants reach high densities




Figure 1. Population growth rate of invasive plant species as a function of reductions in fecundity (seed production or seed viability), shown separately for species with different life forms: (a) monocarpic herbs and grasses, (b) short-lived herbs and grasses, and (c) woody shrubs and trees. The dotted line is shown for a population growth rate of $\lambda = 1$ (see the text for a discussion of population growth rate).




Figure 2. Mean elasticity (see the text for a discussion of elasticity) of the population growth rate to changes in fecundity matrix elements, shown separately for species with different life forms. The error bars represent one standard error. The letters over the bars indicate significant pairwise differences between the groups based on Tukey's HSD (honestly significant difference) test.

where lower vital rates due to density dependence would be expected (Pardini et al. 2009). If density dependence in plant establishment is strong, higher fruit and seed production

will result in higher rates of density-dependence-related death of seedlings and will not result in more established plants. This provides another mechanism by which reductions in seed production and seed viability in cultivars may not result in a less invasive phenotype: Cultivars that produce fewer viable seeds may have recruited offspring similar to those producing more seeds if their offspring suffer less from density-dependent mortality.

In our synthesis, we considered only how reductions in seed production influence the population growth rates of invasive plants. In reality, cultivars may differ in a variety of demographic vital rates (e.g., adult survivorship, seedling growth rates), all of which might affect their invasive potential. For example, the time to reproductive maturity can vary among

cultivars for many species. In *Ardisia crenata*, artificial selection for denser foliage resulted in a cultivar for which individual plants grow more slowly than the wild type. However, this ecotype is not necessarily less invasive, since plants with this denser foliage were found to be stronger competitors for light (Kitajima et al. 2006).

Future research priorities

We are concerned that the marketing of cultivars as “safe to natural areas” has advanced much faster than the research evaluating those cultivars. We recommend two additional lines of study that would advance our ability to evaluate whether a cultivar is likely to threaten natural areas. First, significant demographic differences must exist between cultivars. Studies in which significant differences are found between cultivars in a few vital rates (e.g., the seed production of cultivars) should not be sufficient evidence to justify marketing a cultivar as noninvasive. Our review highlights this point by showing that, in many cases, large reductions in fecundity are inadequate to reduce population growth, particularly for invasive woody plants. Ideally, field trials in which the demography of cultivars is compared would be conducted in all growing zones in which the species is sold. Second, it must be demonstrated that cultivars do not revert to “wild-type” or more weedy forms, either because their offspring do not breed true or through crossing with other cultivars or wild-type individuals. If focal cultivars are capable of hybridizing with other cultivars or wild-type individuals, the fecundity of the offspring may be significantly higher than that of the parent plant. For example, there is evidence that intraspecific hybridization in the

Callery pear (*Pyrus calleryana*), including between different cultivars, can increase invasiveness (Culley and Hardimann 2008). It is possible that some cultivars truly are less invasive than the wild type of the species, and if this is the case, the diversity that can exist among cultivars must be recognized with respect to species bans, and exceptions should be made for those that are highly unlikely to form viable populations in natural areas. However, to date, the research does not exist that would place any nonsterile cultivar in this category.

To our knowledge, these lines of evidence have not been investigated through thorough experimentation on any cultivars of invasive species. An important first step for research is to consider the simplest scenario: If all cultivars of a species except the least fecund one are banned from sale, will the species still pose a significant risk of forming new populations in natural areas and becoming invasive? The answer to this question lies in measuring the vital rates of outcrossed offspring of the cultivar. Only when cultivars do not revert and when their population growth rate is low (close to or less than 1) should they be considered “safe.” We note that completely sterile (female sterile) cultivars can be considered safe without any further investigation. However, we would urge caution in describing self-incompatible cultivars or cultivars that are male but not female sterile as safe. As has been demonstrated in purple loosestrife (*Lythrum salicaria*), these cultivars can set copious seed if they are planted in proximity to other cultivars or feral populations (Anderson and Ascher 1993).

Whenever it is possible, efforts should be focused on creating completely sterile cultivars. Such efforts are underway for some taxa, including *B. thunbergii* and *E. alatus* (Mark Brand, Department of Plant Science and Landscape Architecture, University of Connecticut, Storrs, personal communication, 24 May 2011). Li and colleagues (2004) described a transgenic approach to create “supersterile” cultivars that are both male and female sterile and parthenocarpic. Such molecular approaches to sterility have been applied to many crop plants but, to date, have not been developed for horticultural shrubs. Li and colleagues (2004) argued that supersterile cultivars are highly unlikely to ever reproduce sexually and that the risk of escape of transgenes is greatly reduced as well. Although developing sterility is time consuming and expensive, it provides an alternative to taking a species off the market that it difficult to replace.

Conclusions

We are concerned by the advertisement and marketing of “safe” cultivars of some of the most invasive woody plant species. These claims of their environmental safety are based on very limited demographic evidence that was focused on horticulturally produced plants rather than on their offspring. This article is intended to serve as a thought piece to present this problem to invasive species biologists, as well as to scientists and practitioners in the green industry. Our article highlights that even dramatic changes in some

demographic vital rates—particularly, seed production—may be insufficient to reduce the population growth rate and therefore the invasive potential of a species. Furthermore, offspring of a horticulturally produced plant might look very different from their parent, either because they cross with other cultivars or a wild plant or because they do not breed true. We hope that this article will encourage researchers to consider all demographic vital rates and offspring combinations and will encourage a greater focus on the development of completely sterile plants or on emphasizing native species in horticulture.

Acknowledgments

We thank Jim Ault, Richard Hawke, Ellen Jacquart, and Mark Brand for discussions about cultivars, Amy Ando for discussions about the economic implications of invasive species bans, and Jim Ault and Jon Chase for their comments on the manuscript.

References cited

- Anderson NO, Ascher PD. 1993. Male and female fertility of loosestrife (*Lythrum*) cultivars. *Journal of the American Society for Horticultural Science* 118: 851–858.
- Brown KA, Spector S, Wu W. 2008. Multi-scale analysis of species introductions: Combining landscape and demographic models to improve management decisions about nonnative species. *Journal of Applied Ecology* 45: 1639–1648.
- Buckley YM, Brockerhoff E, Langer L, Ledgard N, North H, Rees M. 2005. Slowing down a pine invasion despite uncertainty in demography and dispersal. *Journal of Applied Ecology* 42: 1020–1030.
- Caswell H. 2001. *Matrix Population Models: Construction, Analysis and Interpretation*, 2nd ed. Sinauer.
- Crone E, et al. 2011. How do plant ecologists use matrix population models? *Ecology Letters* 14: 1–8.
- Culley TM, Hardimann NA. 2008. The role of intraspecific hybridization in the evolution of invasiveness: A case study of the ornamental pear tree *Pyrus calleryana*. *Biological Invasions* 11: 1107–1119.
- Davis AS, Dixon PM, Liebman M. 2004. Using matrix models to determine cropping system effects on annual weed demography. *Ecological Applications* 14: 655–668.
- Davis AS, Landis DA, Nuzzo V, Blosssey B, Gerber E, Hinz HL. 2006. Demographic models inform selection of biocontrol agents for garlic mustard (*Alliaria petiolata*). *Ecological Applications* 16: 2399–2410.
- De Kroon H, Plaisier A, van Groenendael J, Caswell H. 1986. Elasticity: The relative contribution of demographic parameters to population growth rate. *Ecology* 67: 1427–1431.
- DeWalt SJ. 2006. Population dynamics and potential for biological control of an exotic invasive shrub in Hawaiian rainforests. *Biological Invasions* 8: 1145–1158.
- Drew J, Anderson NO, Andow D. 2010. Conundrums of a complex vector for invasive species control: A detailed examination of the horticulture industry. *Biological Invasions* 12: 2837–2851.
- Ehrenfeld JG, Kourtev P, Huang W. 2001. Changes in soil functions following invasions of exotic understory plants in deciduous forests. *Ecological Applications* 11:1287–1300.
- Emery SM, Gross KL. 2005. Effects of timing of prescribed fire on the demography of an invasive plant, spotted knapweed *Centaurea maculosa*. *Journal of Applied Ecology* 42: 60–69.
- Fagan ME, Peart DR. 2004. Impact of the invasive shrub glossy buckthorn (*Rhamnus frangula* L.) on juvenile recruitment by canopy trees. *Forest Ecology and Management* 194: 95–107.

- Frapppier B, Eckert RT, Lee TD. 2004. Experimental removal of the non-indigenous shrub *Rhamnus frangula* (glossy buckthorn): Effects on native herbs and woody seedlings. *Northeastern Naturalist* 11: 333–342.
- Grigulis K, Sheppard AW, Ash JE, Groves RH. 2001. The comparative demography of the pasture weed *Echium plantagineum* between its native and invaded ranges. *Journal of Applied Ecology* 38: 281–290.
- Harrington RA, Kujawski R, Ryan HDP. 2003. Invasive plants and the green industry. *Journal of Arboriculture* 29: 42–48.
- Hyatt LA, Araki S. 2006. Comparative population dynamics of an invading species in its native and novel ranges. *Biological Invasions* 8: 261–275.
- Ingram D, Van Sanford D, Dillon C, Fleming R. 2004. New Crop Opportunities: Phase V. United States Department of Agriculture. (28 July 2011; www.reeis.usda.gov/web/crisprojectpages/199520.html)
- Jacquart EM, Knight TM. 2010. Are there non-invasive cultivars of buckthorn? *Indiana Nursery and Landscape News* 70: 16–17.
- Jacquemyn H, Brys R, Neubert MG. 2005. Fire increases invasive spread of *Molinia caerulea* mainly through changes in demographic parameters. *Ecological Applications* 15: 2097–2108.
- Kitajima K, Fox AM, Sato T, Nagamatsu D. 2006. Cultivar selection prior to introduction may increase invasiveness: Evidence from *Ardisia crenata*. *Biological Invasions* 8: 1471–1482.
- Koop AL, Horvitz CC. 2005. Projection matrix analysis of the demography of an invasive, nonnative shrub (*Ardisia elliptica*). *Ecology* 86: 2661–2672.
- Kourtev PS, Ehrenfeld JG, Huang WZ. 1998. Effects of exotic plant species on soil properties in hardwood forests of New Jersey. *Water, Air, and Soil Pollution* 105: 493–501.
- Lambrecht-McDowell SC, Radosevich SR. 2005. Population demographics and trade-offs to reproduction of an invasive and noninvasive species of *Rubus*. *Biological Invasions* 7: 281–295.
- Lehrer JM, Brand MH, Lubell JD. 2006. Four cultivars of Japanese barberry demonstrate differential reproductive potential under landscape conditions. *HortScience* 41: 762–767.
- Li Y, et al. 2004. Invasive ornamental plants: Problems, challenges and molecular tools to neutralize their invasiveness. *Critical Reviews in Plant Sciences* 23: 381–389.
- Lubell JD, Brand MH, Lehrer JM. 2008. AFLP identification of *Berberis thunbergii* cultivars, inter-specific hybrids and their parental species. *Journal of Horticultural Science and Biotechnology* 83: 55–63.
- Marco DE, Páez SA. 2000. Invasion of *Gleditsia triacanthos* in *Lithraea ternifolia* montane forests of central Argentina. *Environmental Management* 26: 409–419.
- Marinelli J, Hanson B, editors. 1996. *Invasive Plants: Weeds of the Global Garden*. Handbook Number 149. Brooklyn Botanic Garden.
- Nehrbass N, Winkler E, Pergl J, Perglová I, Pyšek P. 2006. Empirical and virtual investigation of the population dynamics of an alien plant under the constraints of local carrying capacity: *Heracleum mantegazzianum* in the Czech Republic. *Perspectives in Plant Ecology, Evolution and Systematics* 7: 253–262.
- Ophardt MC. 2009. Tri-cities shows its fall beauty. *Tri-city Herald*. (28 July 2011; www.tri-cityherald.com/2009/10/24/765571/tri-cities-shows-its-fall-beauty.html)
- Pardini EA, Drake JM, Chase JM, Knight TM. 2009. Complex population dynamics and control of the invasive biennial, *Alliaria petiolata* (garlic mustard). *Ecological Applications* 19: 387–397.
- Parker IM. 2000. Invasion dynamics of *Cytisus scoparius*: A matrix model approach. *Ecological Applications* 10: 726–743.
- Raghu S, Wilson JR, Dhileepan K. 2006. Refining the process of agent selection through understanding plant demography and plant response to herbivory. *Australian Journal of Entomology* 45: 308–316.
- Ramula S, Knight TM, Burns JH, Buckley YM. 2008. General guidelines for invasive plant management based on comparative demography of invasive and native plant populations. *Journal of Applied Ecology* 45: 1124–1133.
- Ranney TG. 2004. Population control: Developing noninvasive nursery crops. *Combined Proceedings of the International Plant Propagators Society* 54: 604–607.
- Reichard SH. 1997. Prevention of invasive plant introductions on national and local levels. Pages 215–227 in Luken JA, Thieret JA, eds. *Assessment and Management of Plant Invasions*. Springer.
- Reichard SH, White P. 2001. Horticulture as a pathway for invasive plant introduction in the United States. *BioScience* 51: 103–113.
- Shea K, Kelly D. 1998. Estimating biocontrol agent impact with matrix models: *Carduus nutans* in New Zealand. *Ecological Applications* 8: 824–832.
- Silvertown J, Franco M, Pisanty I, Mendoza A. 1993. Comparative plant demography: Relative importance of life-cycle components to the finite rate of increase in woody and herbaceous perennials. *Journal of Ecology* 81: 465–476.
- Tenhumberg B, Louda SM, Eckberg JO, Takahashi M. 2008. Monte Carlo analysis for parameter uncertainty in matrix models for weed *Cirsium vulgare*. *Journal of Applied Ecology* 45: 438–447.
- Werner PA, Caswell H. 1977. Population growth rates and age versus stage-distribution models for teasel (*Dipsacus Sylvestris* Huds.). *Ecology* 58: 1103–1111.
- Wilcove DS, Rothstein D, Dubow J, Phillips A, Losos E. 1998. Quantifying threats to imperiled species in the United States. *BioScience* 48: 607–615.
- Williams R. 2004. *Rhamnus* plant named “Ron Williams.” US Patent PP14,791, filed 21 July 2003, and issued 18 May 2004.

Tiffany M. Knight (tknight@wustl.edu) is an associate professor at Washington University in St. Louis, Missouri. Kayri Havens is a director and Pati Vitt is a curator at the Chicago Botanic Garden in Glencoe, Illinois.