

Sivaram Narayan  
Central Michigan University

## Minimal scalings and structural properties of scalable frames

### Abstract

For a unit-norm frame  $F = \{f_i\}_{i=1}^k$  in  $\mathbb{R}^n$ , a scaling is a vector  $c = (c(1), \dots, c(k)) \in \mathbb{R}_{\geq 0}^k$  such that  $\{\sqrt{c(i)}f_i\}_{i=1}^k$  is a Parseval frame in  $\mathbb{R}^n$ . If such a scaling exists,  $F$  is said to be scalable. A scaling  $c$  is a minimal scaling if  $\{f_i : c(i) > 0\}$  has no proper scalable subframe. It is known that the set of all scalings of  $F$  is a convex polytope with vertices corresponding to minimal scalings. In this talk, we provide a method to find a subset of contact points which provides a decomposition of the identity, and an estimate of the number of minimal scalings of a scalable frame. We provide a characterization of when minimal scalings are affinely dependent. Using this characterization, we can conclude that all strict scalings  $c = (c(1), \dots, c(k)) \in \mathbb{R}_{> 0}^k$  of  $F$  have the same structural property. We also present the uniqueness of orthogonal partitioning property of any set of minimal scalings, which provides all possible tight subframes of a given scaled frame

Talk time: 2016-07-21 02:30 PM— 2016-07-21 02:50 PM  
Talk location: Cupples I Room 218