

Washington University in St. Louis

Washington University Open Scholarship

Volume 12

Washington University
Undergraduate Research Digest

Spring 2017

Parkinson's Disease Fibrils: Analyzing Aggregation Growth Properties in a Cell System

María I. Dabrowski

Washington University in St. Louis

Follow this and additional works at: https://openscholarship.wustl.edu/wuurd_vol12

Recommended Citation

Dabrowski, María I., "Parkinson's Disease Fibrils: Analyzing Aggregation Growth Properties in a Cell System" (2017). *Volume 12*. 40.

https://openscholarship.wustl.edu/wuurd_vol12/40

This Abstracts A-I is brought to you for free and open access by the Washington University Undergraduate Research Digest at Washington University Open Scholarship. It has been accepted for inclusion in Volume 12 by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

PARKINSON'S DISEASE FIBRILS:
ANALYZING AGGREGATION GROWTH PROPERTIES
IN A CELL SYSTEM

María I. Dabrowski

Mentors: Rebecca L. Miller and Paul T. Kotzbauer

Parkinson's disease (PD) is the second most common neurodegenerative disease in the world. It occurs when neurons in the substantia nigra degenerate, leading to tremors, rigidity, slow muscle movement and other physical and cognitive impairments. Presently there is no way to accurately diagnose PD prior to a post-mortem autopsy. In approximately less than 1% of PD patients, mutations in alpha-synuclein cause the protein to aggregate. There are five known mutations, and each cause slightly different aggregation outcomes depending on how the proteins misfold and clump together into fibrils. The purpose of our experiment is to generate these mutations in a cell system and test the cells to see how wild-type (WT) and mutated alpha-synuclein-expressing cells cause aggregation of the protein. Doing this in a cell system instead of *in vitro* will allow us to see how cellular lipids and proteins influence monomeric proteins to become integrated into fibrils, and how the aggregation properties differ across mutations. Additionally, completing experiments within a cell system will indicate what natural mechanisms cells have to remove or degrade monomers and fibrils, and how these removal pathways might be impacted by mutations. Studying the varying aggregation morphologies and growth patterns of these mutations will be a step in the right direction toward diagnosing PD and developing better treatments.