

Washington University in St. Louis

Washington University Open Scholarship

Volume 13

Washington University
Undergraduate Research Digest

Spring 2018

Comparative Analysis of Transcription Start Sites and Conserved Motifs in *Drosophila* F and D Elements

Ben French

Washington University in St. Louis

Joe Zielke

Washington University in St. Louis

Follow this and additional works at: https://openscholarship.wustl.edu/wuurd_vol13

Recommended Citation

French, Ben and Zielke, Joe, "Comparative Analysis of Transcription Start Sites and Conserved Motifs in *Drosophila* F and D Elements" (2018). *Volume 13*. 62.

https://openscholarship.wustl.edu/wuurd_vol13/62

This Abstracts A-I is brought to you for free and open access by the Washington University Undergraduate Research Digest at Washington University Open Scholarship. It has been accepted for inclusion in Volume 13 by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

COMPARATIVE ANALYSIS OF TRANSCRIPTION START SITES AND CONSERVED MOTIFS IN *DROSOPHILA* F AND D ELEMENTS

Ben French and Jon Zielke

Mentor: Sarah C. R. Elgin

The *Drosophila melanogaster* Muller F element is unusual in that this chromosome is packaged mostly as heterochromatin, but contains ~80 protein-coding genes. Past studies have shown that classical markers of heterochromatin (e.g., HP1a) are depleted at the transcription start sites (TSS) of active F element genes, which suggests that the factors that regulate F element gene expression are located near the TSS. To define search regions for efforts to identify potential regulatory sites, we manually annotated the TSS positions of genes on the *Drosophila biarmipes* F element (118 TSS) and on a euchromatic region at the base of the D element (258 TSS). These TSS annotations are based on multiple lines of evidence (e.g., sequence homology, RNA-seq data, RNA polymerase II ChIP-seq data). We analyzed promoter shapes (i.e., peaked, intermediate, broad) and the distributions of transcription factor binding sites (TFBS) for these TSS. We found that a substantially greater proportion of D element promoters are classified as peaked compared to F element promoters in both species. To further characterize core promoters of F element genes, we partitioned the *D. biarmipes* F element promoters based on known TFBS observed in the *D. melanogaster* orthologs. We then analyzed each sub-population (partition) using the MEME suite to identify known and novel motifs in the promoters of *D. biarmipes* F element genes. For the dl, twi, Med, da, Udx, zfh1, hb, and Med + dl partitions, no significant motifs (E-value < 1E-05) were found. The most significant motif discovered by MEME was for the zfh1 partition, with an Expect value of 4.3E-04. A Tomtom search of this motif against the FlyFactorSurvey database did not show any significant matches to known TFBS (false discovery rate < 0.01). Hence this motif might be novel.