

Vladimir Bolotnikov

The College of William and Mary

Beurling-Lax type theorems in weighted Bergman-Fock spaces

Abstract

Since the shift operator $M_z : f(z) \rightarrow zf(z)$ is an isometry on the \mathcal{Y} -valued Hardy space $H_{\mathcal{Y}}^2$ of the open unit disk, any M_z -invariant closed subspace $\mathcal{M} \subset H_{\mathcal{Y}}^2$ is generated by the wandering subspace $\mathcal{E} = \mathcal{M} \ominus z\mathcal{M} = P_{\mathcal{M}}z\mathcal{M}^{\perp}$. Furthermore, $z^k\mathcal{E} \perp z^{\ell}\mathcal{E}$ for $k \neq \ell$, and any wandering subspace has the form $\mathcal{E} = \Theta\mathcal{U}$ for some $\mathcal{L}(\mathcal{U}, \mathcal{Y})$ -valued inner function Θ and an appropriate coefficient space \mathcal{U} , which in turn leads to the representations

$$\mathcal{M} = \bigoplus_{k \geq 0} (z^k \mathcal{M} \ominus z^{k+1} \mathcal{M}) = \bigoplus_{k \geq 0} z^k \mathcal{E} = \bigoplus_{k \geq 0} z^k (P_{\mathcal{M}}z\mathcal{M}^{\perp}) = \bigoplus_{k \geq 0} z^k \Theta\mathcal{U} = \Theta H_{\mathcal{U}}^2$$

for an M_z -invariant subspace $\mathcal{M} \subset H_{\mathcal{Y}}^2$. These equivalent representations display the Beurling-Lax theorem and admit extensions to the noncommutative Fock space setting of formal power series in several non-commuting variables. We will discuss their possible extensions in the context of weighted Bergman-Fock spaces where they produce several non-equivalent representations for closed subspaces invariant under multiplication by coordinate functions.

Talk time: 07/19/2016 3:30PM— 07/19/2016 3:50PM

Talk location: Crow 206

Special Session: Multivariable operator theory. Organized by H. Woerdeman.