

Washington University in St. Louis

## Washington University Open Scholarship

---

Volume 12

Washington University  
Undergraduate Research Digest

---

Spring 2017

### Towards Discovering Inhibitors of cyt c Biosynthesis in Systems, I, II, and III

Shannon J. Jiang

*Washington University in St. Louis*

Follow this and additional works at: [https://openscholarship.wustl.edu/wuurd\\_vol12](https://openscholarship.wustl.edu/wuurd_vol12)

---

#### Recommended Citation

Jiang, Shannon J., "Towards Discovering Inhibitors of cyt c Biosynthesis in Systems, I, II, and III" (2017).  
*Volume 12*. 89.

[https://openscholarship.wustl.edu/wuurd\\_vol12/89](https://openscholarship.wustl.edu/wuurd_vol12/89)

This Abstracts J-R is brought to you for free and open access by the Washington University Undergraduate Research Digest at Washington University Open Scholarship. It has been accepted for inclusion in Volume 12 by an authorized administrator of Washington University Open Scholarship. For more information, please contact [digital@wumail.wustl.edu](mailto:digital@wumail.wustl.edu).

# TOWARDS DISCOVERING INHIBITORS OF CYT C BIOSYNTHESIS IN SYSTEMS I, II, AND III

Shannon J. Jiang

*Mentor: Robert Kranz*

Cytochrome c (cyt c) is a heme protein found in most organisms (including human pathogens) that plays an essential role in both aerobic and anaerobic growth. The biogenesis of c-type cytochromes occurs by three different systems (Systems I and II in bacteria and System III in humans). Besides requiring different protein systems, their site of synthesis is also different. Systems I and II function in the periplasmic space while System III functions in the mitochondrial inter-membrane space. These differences may allow for selective targeting of bacterial systems using antimicrobial compounds which may be beneficial in combating infectious bacterial diseases. My goal is to develop a robust assay that detects cyt c maturation quantitatively to allow for subsequent analyses of levels of cyt c maturation in the presence and absence of potential inhibitors. In recombinant *E. coli*, Systems I and II produce cyt c in the periplasm while System III makes cyt c in the cytoplasm. An *in vivo* assay has been developed in the Kranz lab that detects cyt c produced in the periplasm, but cannot detect production in the cytoplasm. Therefore, I have been optimizing an *in vitro* method to extract cyt c from recombinant *E. coli* grown in small volumes (<5mL) and detect it quantitatively by heme stain. Here I present work showing that the *in vitro* assay can detect differences in the amount of cyt c produced within System III. I also explored the optimal conditions for comparing cyt c biogenesis across the three systems in recombinant *E. coli*. Future direction is to test known inhibitors of heme biosynthesis and protein biosynthesis to test the effectiveness of this assay.