

Washington University in St. Louis

Washington University Open Scholarship

Spring 2018

Washington University
Senior Honors Thesis Abstracts

Spring 2018

PI3K Is Required between E14.5-16.5 for the Normal Development of Cochlear Hair Cells

Yutao Su

Washington University in St. Louis

Follow this and additional works at: https://openscholarship.wustl.edu/wushta_spr2018

Recommended Citation

Su, Yutao, "PI3K Is Required between E14.5-16.5 for the Normal Development of Cochlear Hair Cells" (2018). *Spring 2018*. 123.

https://openscholarship.wustl.edu/wushta_spr2018/123

This Abstract for College of Arts & Sciences is brought to you for free and open access by the Washington University

Senior Honors Thesis Abstracts at Washington University Open Scholarship. It has been accepted for inclusion in Spring 2018 by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

PI3K IS REQUIRED BETWEEN E14.5-16.5 FOR THE NORMAL DEVELOPMENT OF COCHLEAR HAIR CELLS

Yutao Su

Mentors: Lu Yang and David M. Ornitz

Hair cells are mechanosensory receptors in the inner ear responsible for transducing oscillations in air pressure into neural signals that are then encoded as sound. Fibroblast growth factor 20 (FGF20) signaling has been implicated in the differentiation of hair cells. Mice lacking a functional copy of *Fgf20* are deaf and have decreased hair cell numbers. However, the mechanisms that regulate differentiation downstream of FGF20 and its receptor, fibroblast growth factor receptor 1 (FGFR1), are not known. There are five major pathways that serve as downstream effectors of FGFR1 in other systems: PLC γ , MAPK (p38), MEK1/2, PI3K, and STAT. But, which one is involved in differentiation in the cochlea? To answer this question, we treated murine cochlear explants with inhibitors that block each major signaling pathways individually during the timeframe when FGFR1 signaling was found to be critical for differentiation, E14.5-E16.5. We found that inhibiting signaling through PLC γ , MEK1/2, and MAPK (p38) did not result in significant changes in total hair cell number. However, treating cochleae with LY294002, a PI3K inhibitor, did result in a decrease in hair cells similar to inhibition of FGFR1 signaling and phenocopies the *in vivo* *Fgf20* knockout phenotype. This suggests that PI3K signaling may play a role in mediating hair cell differentiation. Furthermore, AKT is a common downstream effector of PI3K. Inhibiting AKT, however, showed no decrease in hair cells suggesting that an alternative pathway is responsible for differentiation. We are currently experimenting with alternative signaling pathways downstream of PI3K.