## Gregory Marx

## Virginia Tech

## Completely positive kernels: the noncommutative correspondence setting

## Abstract

It is well known that a function  $K: \Omega \times \Omega \to \mathcal{L}(\mathcal{Y})$  (where  $\mathcal{L}(\mathcal{Y})$  is the set of all bounded linear operators on a Hilbert space  $\mathcal{Y}$ ) being (1) a positive kernel in the sense of Aronszajn (i.e.  $\sum_{i,j=1}^{N} \langle K(\omega_i, \omega_j) y_j, y_i \rangle \geq 0$  for all  $\omega_1, \ldots, \omega_N \in \Omega, y_1, \ldots, y_N \in \mathcal{Y}$ , and  $N = 1, 2, \ldots$ ) is equivalent to (2) K being the reproducing kernel for a reproducing kernel Hilbert space  $\mathcal{H}(K)$ , and (3) K having a Kolmogorov decomposition  $K(\omega, \zeta) = H(\omega)H(\zeta)^*$  for an operator-valued function  $H: \Omega \to \mathcal{L}(\mathcal{X}, \mathcal{Y})$  where  $\mathcal{X}$  is an auxiliary Hilbert space.

Recent work of the authors extended this result to the setting of free noncommutative functions (i.e. functions defined on matrices over a point set which respects direct sums and similarities) with the target set  $\mathcal{L}(\mathcal{Y})$  of K replaced by  $\mathcal{L}(\mathcal{A}, \mathcal{L}(\mathcal{Y}))$  where  $\mathcal{A}$  is a  $C^*$ -algebra. In this talk, we discuss the next extension where the target set of K is replaced by  $\mathcal{L}(\mathcal{A}, \mathcal{L}_a(\mathcal{E}))$  where  $\mathcal{A}$  is a  $W^*$ -algebra and  $\mathcal{L}_a(\mathcal{E})$  is the set of adjointable operators on a Hilbert  $W^*$ -module over a  $W^*$ -algebra  $\mathcal{B}$ . Various special cases of this result correspond to results of Kasparov, Murphy, and Szafraniec in the Hilbert  $C^*$ -module literature.

> Talk time: 07/19/2016 4:00PM— 07/19/2016 4:20PM Talk location: Cupples I Room 113

Special Session: State space methods in operator and function theory. Organized by J. Ball and S. ter Horst.