

Orr Shalit

Technion

## Dilations, inclusions of matrix convex sets, and completely positive maps

### Abstract

In this talk I will present a part of a recent joint work with Davidson, Dor-On, and Solel (a complementary talk will be given by Adam Dor-On in the *Multivariable Operator Theory* special session).

If  $A = (A_1, \dots, A_d)$  is a tuple of operators on  $H$  and  $B = (B_1, \dots, B_d)$  is a tuple of operators on  $K$ , then  $B$  is said to be a *dilation* of  $A$ , denoted  $A \prec B$ , if  $A_i = P_H B_i|_H$  for all  $i$ . For a long time it seemed that the name of the game was: given a commuting tuple of operators  $A$ , find a commuting tuple of *normal* operators  $B$  such that  $A \prec B$  (usually with additional conditions on the joint spectrum  $\sigma(B)$ , and requiring the dilation to hold for powers as well). Quite recently, Helton, Klep, McCullough and Schweighofer changed the rules, and started dilating tuples of *noncommuting* operators to commuting tuples of normal operators. They showed that there is a universal constant  $\vartheta_n$ , such that given a tuple of  $n \times n$  selfadjoint contractions  $A$ , there exists a tuple of commuting selfadjoints  $B$ , such that  $\sigma(B) \subseteq [-1, 1]^d$  and  $\frac{1}{\vartheta_n} A \prec B$ . This result had deep implications to spectrahedral inclusion problems.

The constant  $\vartheta_n$  behaves roughly like  $\sqrt{n}$ , and was shown to be the best constant possible. We were led to ask whether it is possible to obtain such a dilation result with a constant that does not depend on  $n = \text{rank } A$  (necessarily fixing  $d$ ). Moreover, we sought a normal dilation  $B$  with more precise control on the joint spectrum  $\sigma(B)$ . As a representative of our results, I will present the following theorem, as well as some applications.

**Theorem.** *Let  $K$  be a convex set in  $\mathbb{R}^d$  satisfying some reasonable conditions. Then for every  $d$ -tuple  $A$  of selfadjoint operators with a joint numerical range contained in  $K$ , there is a  $d$ -tuple of commuting selfadjoint operators  $B$  with joint spectrum  $\sigma(B) \subseteq K$ , such that*

$$\frac{1}{d} A \prec B.$$

Talk time: 07/20/2016 11:00AM— 07/20/2016 11:30AM

Talk location: Brown Hall 100