

Washington University in St. Louis
Washington University Open Scholarship

Engineering and Applied Science Theses &
Dissertations

Engineering and Applied Science

Spring 2018

The Role of Obesity and Dietary Fatty Acid Content in Regulating Humeral Bone and Cartilage Homeostasis

Lauren Votava

Washington University in St. Louis

Follow this and additional works at: https://openscholarship.wustl.edu/eng_etds



Part of the [Other Biomedical Engineering and Bioengineering Commons](#)

Recommended Citation

Votava, Lauren, "The Role of Obesity and Dietary Fatty Acid Content in Regulating Humeral Bone and Cartilage Homeostasis" (2018). *Engineering and Applied Science Theses & Dissertations*. 353.
https://openscholarship.wustl.edu/eng_etds/353

This Thesis is brought to you for free and open access by the Engineering and Applied Science at Washington University Open Scholarship. It has been accepted for inclusion in Engineering and Applied Science Theses & Dissertations by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

WASHINGTON UNIVERSITY IN ST. LOUIS

School of Engineering & Applied Science
Department of Biomedical Engineering

Dissertation Examination Committee:

Farshid Guilak

Spencer Lake

Amit Pathak

The Role of Obesity and Dietary Fatty Acid Content in
Regulating Humeral Bone and Cartilage Homeostasis
by
Lauren Votava

A thesis presented to
The School of Engineering and Applied Science
of Washington University in
partial fulfillment of the
requirements for the degree
of Master of Science

May 2018
St. Louis, Missouri

© 2018, Lauren Elaine Votava

Table of Contents

List of Figures	iii
Acknowledgments.....	iv
Abstract.....	vi
Chapter 1: Introduction.....	14
Chapter 2: Methods.....	18
2.1 Animal Model	18
2.2 Micro-computed tomographical analysis of trabecular and cortical bone regions	18
2.3 Atomic Force Microscopy.....	19
2.4 Histology	20
2.5 Statistical Analysis	21
Chapter 3: Results.....	22
3.1 Epiphyseal bone microstructure.....	22
3.2 Cortical bone microstructure.....	23
3.3 Atomic Force Microscopy.....	25
2.3 Histology	26
Chapter 4: Discussion.....	28
References.....	32

List of Figures

Figure 3.1: Trabecular region micro-computed tomography results	21
Figure 3.2: Cortical bone micro-computed tomography results	23
Figure 3.3: Atomic force microscopy mechanical mapping results	24
Figure 3.4: Representative histology images	25
Figure 3.5: Histological scoring and thickness results	26

Acknowledgments

I would like to thank Washington University School of Engineering for the incredible opportunities provided to students at this institution as well as our funding sources: NIH R42 AR66439, NIH R42 AR55042, NIH R01 AG46927, NIH R01 AG15768, NIH R21 AR67467, NIH R01 AR48852, NIH R21 AR65956, The Arthritis Foundation, AO Foundation, Department of Defense, NSF EAGER Award, Nancy Taylor Foundation, and Shriners Hospitals for Children.

This project would not have been possible without so many people—only a small number of whom are mentioned here. Dr. Farshid Guilak, as without him, none of this would have been possible. He is an incredible mentor and I've learned so much. Dr. Dennis Barbour for all the time and effort he spent advising me and making this possible. Annie Gitomer-Schwartz, PhD. for teaching me basically every technique I used in this project and then some, for reading over every draft of everything I wrote and every presentation, and so much more. Chia-Lung Wu, PhD. for developing the mice and offering incredible technical support. Natalia Harasymowicz, PhD. and her help with bone microstructure analysis, histology and overall guidance and support. Sara Oswald, M.S. for not only proof-reading a million drafts but also for advice throughout this process. Kelsey Collins, PhD. for dropping everything to grade joints and help with data analysis. Chris Rowland, PhD. for his expert advice on micro-computed tomography, the hours of walking me through how to write a paper, and everything he does to keep the lab running and having fun (science rap anyone?). Nick Thompson, and Jim Maus for all they do every day for all of us, and for the encouragement and support. Lastly I'd like to thank the entire Guilak laboratory. This work is truly a group effort.

Lauren Votava

Washington University in St. Louis

May 2018

Dedicated to my parents and my grandmas,
Jane Votava and Ruth Pienkowski,
for their unwavering support.

Abstract

The Role of Obesity and Dietary Fatty Acid Content in
Regulating Humeral Bone and Cartilage Homeostasis

by

Lauren Votava

Master of Science in Biomedical Engineering

Biomedical Engineering

Washington University in St. Louis, 2018

Professor Farshid Guilak

Objective: The goal of this study was to investigate the effects of different dietary fatty acids in the context of diet-induced obesity on bone and cartilage in the humerus. It is known that obesity increases the severity of injury-induced osteoarthritis in the knee, however it is not fully understood what pathological changes have occurred due to diet alone¹. Additionally, while it is known that shoulder osteoarthritis has a link to obesity, the alterations in this joint are incompletely described.

Methods: In order to examine diet-induced changes in both bone and cartilage, this research utilized mice that had been previously fed diets high in saturated fat (SF), omega-6 fatty polyunsaturated fatty acids (ω -6 PUFA) or omega-3 polyunsaturated fatty acids (ω -3 PUFA) for an earlier study¹. Humeral heads were obtained for testing. Analysis for bone morphometry, bone mineral density, cartilage micro-scale mechanical properties using atomic force microscopy, and histological grading was performed.

Results: Differences in bone morphology and mineral density were seen between diet groups. The high-fat diets in general showed decreased bone quality with the ω -3 PUFA diet being partially protected. Micro-scale cartilage stiffness and overall modified Mankin scores showed no diet-dependence.

Discussion: This study showed that specific types of fatty acids differentially alter bone morphology and mineral density, with no observable changes in the articular cartilage. These findings suggest that in the shoulder, diet-induced obesity by itself may not be a risk factor for osteoarthritis, but may result in other musculoskeletal changes.

Chapter 1: Introduction

Osteoarthritis is a degenerative joint disease that is characterized by the thinning or loss of articulating cartilage surfaces, as well as altered bone remodeling and joint pain during movement². There are currently no disease-modifying pharmacological treatments available for osteoarthritis (OA), and the standard of care is to prescribe painkillers until a total joint replacement is required³. One of the primary risk factors for OA is obesity; however, little research had been performed on this topic until recently, as historically OA was considered to be a disease of “wear and tear” or mechanical overloading. Patient data, including the link between obesity and hand and shoulder OA indicates that non-weight-bearing joints can be affected as well. Investigating this topic further, recent research has revealed that metabolic and inflammatory factors, including the content of dietary fatty acids, can significantly interact with mechanical factors to influence the pathogenesis of OA^{1,4}.

Pathogenesis of OA can affect more than just the articulating surface of cartilage. Increasingly this disease is known to be an whole-joint disease, and alterations in bone remodeling, bone mineral content, and synovitis, are known to occur⁵. In some studies, these changes have been shown to precede cartilage degradation markers of OA⁵, although different types of OA may be initiated by differing mechanisms. While the precise timing of the bone changes leading to joint degradation have yet to be elucidated, several animal models of OA show an increase in subchondral trabecular separation and a decrease in bone volume fraction and mineral content within the affected joint^{2,6}. Ultimately, these factors vary during the course of OA progression, but their direct influence on OA remains unknown. Furthermore, the biological factors associated

with obesity have also been found to contribute to bone formation and remodeling. Traditionally, the increased mechanical loading due to excess weight was thought to have an anabolic effect on bone, but studies have shown a decrease in overall bone quality with obesity⁷. This suggests that the increased incidence of OA in obese patients is not simply the result of increased mechanical forces.

Like bone, articular cartilage is sensitive to both mechanical and biological factors. Biological signals such as inflammatory cytokines can lead to upregulation of catabolic factors like matrix metalloproteinase (MMP) activity and decreases in anabolic cues like basic fibroblast growth factor (FGF)⁸. Both increased MMP activity and a decrease in FGF concentration point to amplified catabolic activity and, when accompanied by an upregulation of ADAMTS (a disintegrin and metalloproteinase with thrombospondin motifs), induce hallmarks of OA-like chondrocyte hypertrophy, osteocyte formation and cartilage matrix degeneration⁸⁻¹⁰. This degeneration can be seen on the microscale using atomic force microscopy to map the matrix stiffness in not only the bulk matrix, but also regions surrounding chondrocytes. Chondrocytes are surrounded by a region known as the pericellular matrix (PCM), which differs both in composition and mechanical properties to the bulk extracellular matrix (ECM). Previous studies have seen decreases in the stiffness of the PCM and the ECM with OA, as well as a decrease in the stiffness gradient between the softer PCM and stiffer ECM¹¹. Decreases in the matrix stiffness would alter the mechanical environment of chondrocytes and are hypothesized to increase the forces sensed by the cell, thus leading to release of inflammatory mediators and more degeneration¹².

The complex interplay between biological signals and mechanical changes that occurs with OA is further complicated in obesity. Obesity is the result of excess adipose tissue that is now known to

be metabolically active and leads to systemic inflammation⁴. White adipose tissue releases both adipokines and cytokines that are implicated in OA progression⁴. The combination of obesity-induced pathophysiological loading and systemic inflammation from excess white adipose tissue can lead to cartilage degeneration, bone remodeling, and eventually symptomatic OA in multiple sites including the shoulder^{8,13,14}. Research has shown correlations between adipokines upregulated in obesity and shoulder OA, but little is known about the link beyond observational studies¹⁵. To study the relationship between obesity and OA, animal models of obesity have been generated by feeding a high-fat diet^{1,14,16-18}. Multiple studies have shown increased severity of surgically-induced OA for animals fed a high-fat diet, and the composition of the diet affects the severity of the disease^{1,14}. High-fat diets composed of mainly saturated fatty acids (SFA) or omega-6 poly-unsaturated fatty acids (ω -6 PUFA) show considerably worse knee OA than weight-matched counterparts on a high-fat diet with omega-3 poly-unsaturated fatty acids (ω -3 PUFA) or a normal mouse chow diet¹. These studies show that diets rich in ω -3 protect the knee from injury-induced OA while saturated fat and ω -6 PUFA seem to predispose the joint to increased damage after trauma.

While previous studies have shown OA progression to be dependent on dietary fatty acids after trauma to the joint, it is not fully understood whether diet-induced obesity causes OA in joints other than the knee. The presence of pathological changes in joint structure and stiffness due to diet alone may help to explain the increased degradation seen in certain types of diets and the protective effects of others. Additionally, most studies focus on knee or hip OA in the context of obesity, yet it is known that adipokine receptors are upregulated in chondrocytes from osteoarthritic shoulders and that adipokine concentrations in the joint correlate with body mass index (BMI)¹⁵. Shoulders are also the next most common site of OA after knees, with a link

between knee pain and shoulder pain¹⁹. Clinically, however, there is not a strong association between obesity and OA of the shoulder²⁰. The goal of this investigation is to determine whether alterations in bone structure and cartilage integrity can be seen in the absence of trauma-induced OA to better understand the link between diet, obesity and predisposition to post-traumatic OA.

Chapter 2: Methods

2.1 Animal Model

All animal use procedures were approved by the local IACUC. Mice (C57BL/6) were fed diets high in saturated fat, omega-6 poly-unsaturated fatty acids (ω -6 PUFA), omega-3 poly-unsaturated fatty acids (ω -3 PUFA) or normal mouse chow for 24 weeks beginning at 4 weeks of age¹. Initial reports on knee joints and full diet information can be found in Wu et al. 2014¹. All animal specimens were stored at -20°C following euthanasia. Prior to analysis, specimens were thawed at 4°C, and humeral heads were isolated.

2.2 Micro-computed tomographical analysis of trabecular and cortical bone regions

Humeral heads were fixed in 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield PA) for 24 hours at room temperature. They were then dehydrated in ethanol and scanned in air using microCT at 9 μ m/pixel resolution, and an x-ray voltage of 55kV (Bruker SkyScan 1176). Trabecular regions began at the end of the subchondral plate and ended at the growth plate. These regions were thresholded and analyzed using the BoneJ extension in ImageJ²¹. Bone mineral density was calculated in both trabecular and cortical regions and was calibrated using hydroxyapatite phantoms obtained from Bruker. Trabecular bone volume fraction and trabecular thickness and separation, were calculated (n=12-15). Cortical bone regions were analyzed using the automated CTan software package (Bruker). Cortical bone cross-sectional area and thickness were calculated in two diaphyseal regions one 0.5mm from the humeral head and another at the end of the deltoid tuberosity (n=6-14).

2.3 Atomic Force Microscopy

Mechanical testing with an atomic force microscope was performed as described previously^{22,23}.

In brief, freshly dissected humeral heads were embedded in optimum cutting temperature (O.C.T. Tissue-Tek, Sakura Finetek USA, Inc. Torrance, CA) medium and 5 μ m thick cryosections of humeral head articular cartilage were obtained. Sections were briefly thawed and immunologically stained for collagen IV due to its exclusive presence in the PCM (Affinity Purified Rabbit Polyclonal Collagen Type VI Antibody; Fitzgerald Industries, Acton, MA, Goat Anti-Rabbit Alexa Fluor® 488; abcam, Cambridge, MA). Briefly, sections were washed in PBS, blocked in 10% normal goat serum (Life Technologies, Frederick, MD), and incubated with the primary antibody diluted 1:200 in blocking solution for 1 hour. Sections were rinsed with PBS and stained with Alexa-fluor488 conjugated goat anti rabbit antibody diluted at 1:200. Sections were maintained in PBS until analysis to preserve hydration. The local mechanical modulus was calculated using an atomic force microscope (MFP-3D, Asylum Research, Santa Barbara, CA) cantilever with a 5 μ m-diameter borosilicate spherical indenter ($k = 5.4\text{N/m}$; Novascan Technologies, Ames, IA) that was calibrated daily. Regions of interest were selected approximately 1-2 cells away from the outer edges of the sections to avoid edge artifacts while remaining out of deep-zone cartilage and subchondral bone. Each region was 10 μm^2 and an indentation was performed every 0.5 μm to give a total number of 400 indentations per region. The sample was indented at a rate of 10 $\mu\text{m/s}$ as previous studies have shown little or no variation in modulus values for indentation speeds between 5 and 25 $\mu\text{m/s}$ ²⁴. The indentation was continued until the trigger force, 300nN, was reached.

Cantilever deflection and z-piezo movement were obtained from the Asylum Research software and analyzed using a custom MATLAB script (The MathWorks, Natick, MA). For determination

of the elastic modulus, a modified Hertz model was utilized as it has been previously applied to atomic force microscope (AFM) analysis of cartilaginous tissue^{22,24-27}. Contact point extrapolation was used to determine the point at which the cantilever contacted the surface as previously described²⁸. A threshold was applied to the fluorescent images to be used as an overlay for the stiffness maps to assign modulus values to either ECM or PCM. Analysis points within the cell body were excluded from further analysis. Data points were excluded if they exceeded 2.5 times the surrounding values and were replaced with the mean of the average value of the adjacent points. To plot the radially changing modulus, the cell region was thresholded in MATLAB and average moduli in increasing 0.5 μ m thick rings out from the edge of the cell were calculated. Due to the close proximity of other chondrocytes in murine cartilage, the data was truncated at its maximum value to avoid including the edges of PCM from the neighboring chondrocytes.

2.4 Histology

Humeral heads were fixed in 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield PA), decalcified using Calce II (Fisher) and embedded in paraffin. The samples were sectioned at 8 μ m and stained using safranin-O, fast green and hematoxylin. The cartilage phenotype was graded using established procedures for OA grading including a modified Mankin scoring system. This consisted of the following categories: cartilage surface structure (0-11), tidemark duplication (0-3), Safranin-O staining (0-8), chondrocyte clones in uncalcified cartilage (0-2), hypertrophic chondrocytes in calcified cartilage (0-2), and subchondral bone thickness (0-2). The total score possible was 28.


2.5 Statistical Analysis

Significance between diet groups for bone and AFM outcomes were determined by one-way ANOVA and Tukey post-hoc tests. For discrete histological grading scores, significance was determined using a Kruskal-Wallis test. Data is shown as mean \pm standard deviation. Outliers were determined using a ROUT test (Q=1%). All statistics were performed in GraphPad Prism 7.03.

Chapter 3: Results

3.1 Epiphyseal bone microstructure

Micro-computed tomography (microCT) analysis of the humeral epiphyseal trabecular region indicated in Figure 3.1A showed no changes in the bone mineral density of the trabeculae (Figure 3.1B) or the trabecular thickness (Figure 3.1D). Bone volume fraction (Figure 3.1C) and trabecular separation (Figure 3.1E) were significantly different between diet groups. Bone volume fraction was significantly decreased in both the ω -6 PUFA and saturated fat diet groups whereas trabecular separation increased for the saturated fat diet group alone compared to control diet group.



3.2 Cortical bone microstructure

In the cortical bone, two regions were analyzed and are shown as the upper cortical region and lower cortical region in Figure 3.2A. More pronounced changes in the bone mineral density can be seen in the lower cortical region, but significant differences due to diet are seen in both (Figure 3.2B and 3.2C). Interestingly, the saturated fat diet shows a lower bone mineral density in the upper region, but not in the lower region. Figure 3.2D and 3.2E show the average total cross-sectional area for each diet group. In the lower region, all high-fat diets result in a decrease in the total cross-sectional area, however in the upper region, only the ω -3 PUFA diet group had a decrease in area. The upper region cortical thickness in the ω -3 PUFA group was also decreased from all other diets, but the lower region thickness showed no diet-dependence.




Figure 3.2: Cortical bone micro-computed tomography results. A) Visual of the two diaphyseal regions analyzed. The upper region begins 0.5mm below the end of the humeral head and is 0.5mm thick. The lower region begins after the deltoid tuberosity and is 0.4mm thick. B) Upper cortical region bone mineral density. The ω-6 PUFA and saturated fat diet groups were significantly different from the control diet, and the saturated fat diet was significantly different from the ω-3 PUFA diet as well. C) Lower cortical region bone mineral density. Contrary to what was seen in the upper region, the ω-6 PUFA diet group showed an increase in mineral density in this region as compared to the other diet groups. D) The upper cortical region cross-sectional area. A significant decrease in the cross-sectional area is seen in the ω-3 PUFA diet group. E) Lower cortical region cross-sectional area. Decreases in the cross-sectional area are seen for all high-fat diets as compared to control. F) Upper cortical region thickness. The ω-3 PUFA diet group showed a decrease in the cortical thickness as compared to all other diets. G) Lower cortical region thickness. No significant differences were seen between diet groups. All statistical significance was determined using a two-way ANOVA. $p < 0.05$, different letters are significantly different from each other. All data is presented as mean \pm standard error of the mean.

3.3 Atomic Force Microscopy

Immunofluorescence-guided atomic force microscopy mechanical testing showed no changes in the modulus values either in the bulk ECM (Figure 3.3A) or the PCM surrounding the cell (Figure 3.3B). The progression of the modulus from softer PCM to stiffer ECM was unchanged due to diet (Figure 3.3C). An example stiffness map is shown in Figure 3.3D. This map shows the softer PCM surrounding the cell and the gradual progression out to stiffer ECM.





Figure 3.3: Atomic force microscopy mechanical mapping results. A) Average ECM Young's modulus. No significant differences were seen due to diet. B) Average PCM Young's modulus. No significant differences were seen due to diet. C) Young's modulus progression out from the cell in concentric rings of 0.5 μ m. High variability between mice was seen and no significant difference in slope due to diet was observed. D) An example stiffness map. Each pixel represents one force plot for a total of 400 indentations for each 10 μ m² area. Statistical significance was tested using a two-way ANOVA. All data is shown as mean \pm standard error of the mean.


2.3 Histology

Example histology images are shown in Figure 3.4. A full humeral head is shown in Figure 3.4A and enlarged images of the cartilage surface are shown in Figure 3.4B. Full modified Mankin score results are shown in Figure 3.5.



While an overall Mankin score showed no differences with diet (Figure 3.5A), two sub-scores that form part of the Mankin scoring system varied with diet. The number of hypertrophic chondrocytes in calcified cartilage (Figure 3.5B) was increased in the ω -6 PUFA and saturated fat groups and the number of chondrocyte clones in uncalcified cartilage (Figure 3.5C) was decreased in the ω -6 PUFA diet. Corroborating the absent cartilage phenotype, the thickness of the uncalcified cartilage (Figure 3.5D) was independent of diet, while the thickness of the calcified cartilage shows statistically insignificant trends (Figure 3.5E). Possibly due to the same

mechanisms that caused the decrease in cortical thickness in Figure 3.2F, the thickness of the subchondral plate was significantly decreased in the ω -3 PUFA diet (Figure 3.5F).



Chapter 4: Discussion

This study showed that a high-fat diet in the absence of joint injury was sufficient to cause detrimental changes in bone remodeling in the humerus both in the trabecular region as well as the cortical region. The changes seen involved alterations both in bone mineral density as well as structural changes that have been linked to an increase in fracture risk like cortical area. Unlike bone, the properties of the cartilage were mostly unaffected by a high-fat diet, even on the microscale. Matrix stiffness and the gradient between softer PCM and stiffer ECM were unchanged. Both stiffness and gradient values have been shown to decrease with OA, and the lack of a statistically significant difference in stiffness measurements on this length scale indicate the relative mechanical integrity of the cartilage. Corroborating the AFM findings, no significant differences in histological scoring for traditional signs of cartilage degeneration were found. Two subscores from the modified Mankin scoring system showed significant differences between diet groups: the number of hypertrophic chondrocytes in the calcified cartilage and the number of chondrocyte clones in the uncalcified region, but the cartilage overall showed few signs of alterations. Another histological measurement, the average thickness of the subchondral plate, was significantly reduced in the ω -3 PUFA diet group compared to the ω -6 PUFA diet and saturated fat diet.

Similar to the controversy that has surrounded the impact of obesity on OA, the effect of obesity on bone quality is currently under investigation. An increase in body weight is expected to lead to an increase in bone strength to support the extra weight during daily activities²⁹. However obesity has been linked to a site-specific increase in fracture risk, including an increased risk in the proximal humerus³⁰. This risk has been partially attributed to a decrease in mineral content, but the relationship between obesity and bone mineral density is incompletely understood. Some

clinical studies indicate a positive relationship between body mass index (BMI) and bone mineral density (BMD), but others show the opposite or no change²⁹. These differences are due in part to the differences in age, sex, genetics, and selection of bone region to analyze.

Differences are also seen in animal models of obesity with some studies showing an increase in bone formation and others a decrease^{31,32}. Our results correlate with a study by Cao et al. using the same mouse strain and a diet slightly lower in fat (45% as compared with 60%) which saw a decrease in cancellous bone mass but not cortical bone mass in the tibia³³. This previous study showed a decrease in trabecular bone volume fraction as well as trabecular separation as was seen in our data³³. However Cao et al. show no significant effects of high-fat feeding on tibial cortical thickness or total area³³. Differences in the diet composition, age of the mice, length of the study and tibia versus humerus could all account for the variation in outcomes. It has been hypothesized that obesity initially leads to overall bone formation due to the increase in body weight, but eventually the systemic inflammation results in a decrease in bone formation³⁴. This process has been hypothesized to occur at different rates in different bones, stressing the need for further investigation into the response of humeral bone to obesity³⁵.

While obesity has clearly been shown to affect bone, the type of fat present in the diet has also been shown to alter the response of bone to a high-fat diet. ω -3 PUFA have been associated with positive cardiovascular health outcomes among others, but a study found a decrease in cortical area and maximal load with high levels of supplementation in developing rabbits³⁶. This decrease in cortical area agrees with our results in the upper cortical region. The decrease in bone volume for the ω -3 PUFA diet has been hypothesized to be caused by the upregulation of adiponectin in these mice as reported previously¹. Adiponectin, acting through FoxO1 has been shown in some situations to decrease osteoblast proliferation and promote apoptosis³⁷. Under what conditions ω -

3 PUFAs result in catabolic or anabolic effects is still an active area of research, but many studies report protective effects of ω -3 PUFA supplementation on bone²⁹. Although awareness of ω -3 PUFA benefits is increasing, the average American diet has proportionally more ω -6 PUFA and saturated fats³⁸. The relationship between ω -6 PUFA and bone is under investigation, but prostaglandin E₂, a downstream metabolite of a prevalent ω -6 PUFA, is linked to an increase in inflammation and may inhibit bone formation at high doses of ω -6 PUFA^{31,35}. Intermediate doses of ω -6 PUFA show conflicting results, with clinical studies finding a positive correlation between ω -6 PUFA consumption and overall BMD in a study of post-menopausal women³⁹. Saturated fatty acid intake does not show the conflicting results that ω -6 PUFA consumption does, with most but not all studies concluding that saturated fats lead to detrimental bone changes^{31,32}. An *in vitro* study showed that saturated fat led to an increase in osteoclast survival and an *in vivo* study showed a decrease in bone volume fraction and an increase in trabecular separation after a high-fat diet^{40,41}. These *in vivo* results correspond with our findings in this study.

The effects of a high-fat diet on cartilage and OA are also investigated in the literature. A series of studies showed that dietary supplementation with lard altered the progression but not the incidence of OA in mice^{42,43}. Other studies have determined an increase in OA score due to a high-fat diet alone, while other studies saw no difference in Mankin scores between a chow diet and a high-fat diet^{17,44}. This study found no difference in overall OA scores due to diet alone, indicating that, at least in the shoulder, bone changes occur before cartilage degeneration or that cartilage degeneration requires an initial trauma to the joint. This was corroborated by the lack of microscale cartilage stiffness changes that are known to occur with OA¹¹. The lack of macro- or microscale cartilage alterations could be due to a number of factors. Changes may occur on a

longer time scale than 14 weeks on a high fat diet, and this time scale as well as the degree of damage may be joint-specific.

In this study, we found diet-specific changes in bone due to obesity in the absence of trauma, while the cartilage surface remained relatively unchanged. Studies have shown joint-specific differences in response to obesity-related OA, with correlations being seen in knee and hand OA but with little or no correlation for hip OA^{4,15}. Literature on shoulder OA is relatively sparse, and future work may shed more light on the pathogenesis of shoulder OA as well as the link between it and obesity.

References

1. Wu, C. *et al.* Dietary fatty acid content regulates wound repair and the pathogenesis of osteoarthritis following joint injury. *Ann. Rheum. Dis.* 1–8 (2015).
doi:10.1136/annrheumdis-2014-205601
2. Man, G. S. & Mologhianu, G. Osteoarthritis pathogenesis - a complex process that involves the entire joint. *J. Med. Life* **7**, 37–41 (2014).
3. Bottini, M. *et al.* Nanodrugs to target articular cartilage: An emerging platform for osteoarthritis therapy. *Nanomedicine Nanotechnology, Biol. Med.* **12**, 255–268 (2016).
4. Aspden, R. M. Obesity punches above its weight in osteoarthritis. *Nat. Rev. Rheumatol.* **7**, 65–68 (2011).
5. Baker-LePain, J. C. & Lane, N. E. Role of bone architecture and anatomy in osteoarthritis. *Bone* **51**, 197–203 (2012).
6. Li, G. *et al.* Subchondral bone in osteoarthritis: insight into risk factors and microstructural changes. *Arthritis Res. Ther.* **15**, 223 (2013).
7. Shapses, S. A. & Sukumar, D. Bone metabolism in obesity and weight loss. *Annu. Rev. Nutr.* **32**, 287–309 (2012).
8. Scotece, M. & Mobasheri, A. Leptin in osteoarthritis: Focus on articular cartilage and chondrocytes. *Life Sci.* **140**, 75–78 (2015).
9. Hamada, D. *et al.* Suppressive Effects of Insulin on Tumor Necrosis Factor-Dependent Early Osteoarthritic Changes Associated With Obesity and Type 2 Diabetes Mellitus. *Arthritis Rheumatol. (Hoboken, N.J.)* **68**, 1392–402 (2016).
10. Wang, X., Hunter, D., Xu, J. & Ding, C. Metabolic triggered inflammation in osteoarthritis. *Osteoarthr. Cartil.* **23**, 22–30 (2015).
11. Wilusz, R. E., Zauscher, S. & Guilak, F. Micromechanical mapping of early osteoarthritic changes in the pericellular matrix of human articular cartilage. *Osteoarthr. Cartil.* **21**, 1895–1903 (2013).
12. Anderson, D. D. *et al.* Post-traumatic osteoarthritis: Improved understanding and opportunities for early intervention. *J. Orthop. Res.* **29**, 802–809 (2011).
13. Courties, A., Gualillo, O., Berenbaum, F. & Sellam, J. Metabolic stress-induced joint inflammation and osteoarthritis. *Osteoarthr. Cartil.* **23**, 1955–1965 (2015).
14. Brunner, A. M. *et al.* High dietary fat and the development of osteoarthritis in a rabbit model. *Osteoarthr. Cartil.* **20**, 584–592 (2012).

15. Gandhi, R., Takahashi, M., Rizek, R., Dessouki, O. & Mahomed, N. N. Obesity-related adipokines and shoulder osteoarthritis. *J. Rheumatol.* **39**, 2046–8 (2012).
16. Collins, K. H. *et al.* Relationship between inflammation, the gut microbiota, and metabolic osteoarthritis development: studies in a rat model. *Osteoarthr. Cartil.* **23**, 1989–1998 (2015).
17. Louer, C. R. *et al.* Diet-induced obesity significantly increases the severity of posttraumatic arthritis in mice. *Arthritis Rheum.* **64**, 3220–3230 (2012).
18. Mooney, R. A., Sampson, E. R., Lerea, J., Rosier, R. N. & Zuscik, M. J. High-fat diet accelerates progression of osteoarthritis after meniscal/ligamentous injury. *Arthritis Res. Ther.* **13**, R198 (2011).
19. Laslett, L. L., Otahal, P., Hensor, E. M. A., Kingsbury, S. R. & Conaghan, P. G. Knee Pain Predicts Subsequent Shoulder Pain and the Association Is Mediated by Leg Weakness: Longitudinal Observational Data from the Osteoarthritis Initiative. *J. Rheumatol.* **43**, 2049–2055 (2016).
20. Cho, H. J., Morey, V., Kang, J. Y., Kim, K. W. & Kim, T. K. Prevalence and Risk Factors of Spine, Shoulder, Hand, Hip, and Knee Osteoarthritis in Community-dwelling Koreans Older Than Age 65 Years. *Clin. Orthop. Relat. Res.* **473**, 3307–14 (2015).
21. Doube, M. *et al.* BoneJ: Free and extensible bone image analysis in ImageJ. *Bone* **47**, 1076–1079 (2010).
22. Wilusz, R. E., Zauscher, S. & Guilak, F. Micromechanical mapping of early osteoarthritic changes in the pericellular matrix of human articular cartilage. *Osteoarthr. Cartil.* **21**, 1895–1903 (2013).
23. Wilusz, R. E., DeFrate, L. E. & Guilak, F. Immunofluorescence-guided atomic force microscopy to measure the micromechanical properties of the pericellular matrix of porcine articular cartilage. *J. R. Soc. Interface* **9**, (2012).
24. Zelenski, N. A. *et al.* Collagen VI regulates pericellular matrix properties, chondrocyte swelling, and mechanotransduction in articular cartilage HHS Public Access. *Arthritis Rheumatol* **67**, 1286–1294 (2015).
25. Harding, J. W. & Sneddon, I. N. The elastic stresses produced by the indentation of the plane surface of a semi-infinite elastic solid by a rigid punch. *Math. Proc. Cambridge Philos. Soc.* **41**, 16 (1945).
26. Darling, E. M., Zauscher, S. & Guilak, F. Viscoelastic properties of zonal articular chondrocytes measured by atomic force microscopy. *Osteoarthr. Cartil.* **14**, 571–579 (2006).

27. Darling, E. M., Wilusz, R. E., Bolognesi, M. P., Zauscher, S. & Guilak, F. Spatial Mapping of the Biomechanical Properties of the Pericellular Matrix of Articular Cartilage Measured In Situ via Atomic Force Microscopy. *Biophys. J.* **98**, 2848–2856 (2010).
28. Darling, E. M., Topel, M., Zauscher, S., Vail, T. P. & Guilak, F. Viscoelastic properties of human mesenchymally-derived stem cells and primary osteoblasts, chondrocytes, and adipocytes. *J. Biomech.* **41**, 454–464 (2008).
29. Shapses, S. A., Pop, L. C. & Wang, Y. Obesity is a concern for bone health with aging. *Nutr. Res.* **39**, 1–13 (2017).
30. Walsh, J. S. & Vilaca, T. Obesity, Type 2 Diabetes and Bone in Adults. *Calcif. Tissue Int.* **100**, 528–535 (2017).
31. Lau, B. Y. *et al.* Influence of high-fat diet from differential dietary sources on bone mineral density, bone strength, and bone fatty acid composition in rats. *Appl. Physiol. Nutr. Metab.* **35**, 598–606 (2010).
32. Wang, Y. *et al.* High fat diet enriched with saturated, but not monounsaturated fatty acids adversely affects femur, and both diets increase calcium absorption in older female mice. *Nutr. Res.* **36**, 742–750 (2016).
33. Cao, J. J., Gregoire, B. R. & Gao, H. High-fat diet decreases cancellous bone mass but has no effect on cortical bone mass in the tibia in mice. *Bone* **44**, 1097–1104 (2009).
34. Lecka-Czernik, B., Stechschulte, L. A., Czernik, P. J. & Dowling, A. R. High bone mass in adult mice with diet-induced obesity results from a combination of initial increase in bone mass followed by attenuation in bone formation; implications for high bone mass and decreased bone quality in obesity. *Mol. Cell. Endocrinol.* **410**, 35–41 (2015).
35. Corwin, R. L. Effects of dietary fats on bone health in advanced age. *Prostaglandins, Leukot. Essent. Fat. Acids* **68**, 379–386 (2003).
36. Judex, S. *et al.* Dietary Fish Oil Supplementation Adversely Affects Cortical Bone Morphology and Biomechanics in Growing Rabbits. *Calcif. Tissue Int.* **66**, 443–448 (2000).
37. Kajimura, D. *et al.* Adiponectin regulates bone mass via opposite central and peripheral mechanisms through FoxO1. *Cell Metab.* **17**, 901–15 (2013).
38. Enos, R. T. *et al.* Reducing the Dietary Omega-6:Omega-3 Utilizing α -Linolenic Acid; Not a Sufficient Therapy for Attenuating High-Fat-Diet-Induced Obesity Development Nor Related Detrimental Metabolic and Adipose Tissue Inflammatory Outcomes. *PLoS One* **9**, e94897 (2014).
39. Järvinen, R. *et al.* Associations of dietary polyunsaturated fatty acids with bone mineral

- density in elderly women. *Eur. J. Clin. Nutr.* **66**, 496–503 (2012).
40. Oh, S.-R. *et al.* Saturated fatty acids enhance osteoclast survival. *J. Lipid Res.* **51**, 892–9 (2010).
 41. Cao, J. J., Sun, L. & Gao, H. Diet-induced obesity alters bone remodeling leading to decreased femoral trabecular bone mass in mice. *Ann. N.Y. Acad. Sci* doi:10.1111/j.1749-6632.2009.05252.x
 42. Silberberg, M. & Silberberg, R. Age Factor and High-Fat Diets in the Evolution of Osteoarthritis in Mice. *J. Gerontol.* **12**, 9–13 (1957).
 43. Griffin, T. M. & Guilak, F. Why is obesity associated with osteoarthritis? Insights from mouse models of obesity. *Biorheology* **45**, 387–98 (2008).
 44. Griffin, T. M., Huebner, J. L., Kraus, V. B., Yan, Z. & Guilak, F. Induction of osteoarthritis and metabolic inflammation by a very high-fat diet in mice: Effects of short-term exercise. *Arthritis Rheum.* **64**, 443–453 (2012).