

Raul Curto

University of Iowa

An algorithmic approach to the sextic moment problem

Abstract

We describe an approach to solving the sextic moment problem based on a thorough analysis of the extremal case, and algorithmic solutions for the non-extremal cases.

For a degree $2n$ complex sequence $\gamma \equiv \gamma^{(2n)} = \{\gamma_{ij}\}_{i,j \in \mathbb{Z}_+, i+j \leq 2n}$ to have a representing measure μ , it is necessary for the associated moment matrix $M(n)$ to be positive semidefinite, and for the algebraic variety associated to γ , $\mathcal{V}_\gamma \equiv \mathcal{V}(M(n))$, to satisfy $\text{rank } M(n) \leq \text{card } \mathcal{V}_\gamma$ as well as the following consistency condition: if a polynomial $p(z, \bar{z}) \equiv \sum_{ij} a_{ij} \bar{z}^i z^j$ of degree at most $2n$ vanishes on \mathcal{V}_γ , then the *Riesz functional* $\Lambda(p) \equiv p(\gamma) := \sum_{ij} a_{ij} \gamma_{ij} = 0$.

Positive semidefiniteness, recursiveness, and the variety condition of a moment matrix are necessary and sufficient conditions to solve the quadratic ($n = 1$) and quartic ($n = 2$) moment problems. Also, positive semidefiniteness, combined with the above mentioned consistency condition, is a sufficient condition in the case of *extremal* moment problems, i.e., when the rank of the moment matrix (denoted by r) and the cardinality of the associated algebraic variety (denoted by v) are equal. However, these conditions are not sufficient for *non-extremal* (i.e., $r < v$) sextic ($n = 3$) or higher-order truncated moment problems.

DTLpar For extremal sextic moment problems, verifying consistency amounts to having good representation theorems for sextic polynomials in two variables vanishing on the algebraic variety of the moment sequence. We obtain such representation theorems using the Division Algorithm from algebraic geometry. As a consequence, we are able to complete the analysis of extremal sextic moment problems.

Assume now that $M(3) \geq 0$, and that it satisfies the variety condition $r \leq v$ as well as consistency. Also assume that $M(3)$ admits at least one *cubic* column relation. We prove the existence of a related matrix $\widetilde{M}(3)$ with $\text{rank } \widetilde{M}(3) < \text{rank } M(3)$ and such that each representing measure for $\widetilde{M}(3)$ gives rise to a representing measure for $M(3)$. As a concrete application, we discuss the case when $\text{rank } M(3) = 8$ and $\text{card } \mathcal{V}(M(3)) \leq 9$.

Along the way, we settle three key instances of the non-extremal sextic moment problem, as follows: when $r = 7$, positive semidefiniteness, consistency and the variety condition guarantee the existence of a 7-atomic representing measure; when $r = 8$ we construct two determining algorithms, corresponding to the cases $v = 9$ and $v = +\infty$. To accomplish this, we generalize the above mentioned rank-reduction technique, which was used in previous work to find an explicit solution of the nonsingular quartic moment problem.

The talk is based on joint work with Seonguk Yoo.

Talk time: 7/21/2016 2:30PM— 7/21/2016 2:50PM

Talk location: Cupples I Room 113

Special Session: Finite and infinite dimensional moment problems. Organized by M. Infusino, S. Kuhlmann, and T. Kuna.